

Helminth Parasites of *Clarias gariepinus* (Clariidae) in Lekki Lagoon, Lagos, Nigeria

Akinsanya B. & O.A. Otubanjo

Department of Zoology, Parasitology Unit, University of Lagos, Lagos, Nigeria.

Corresponding author: Akinsanya, B.: akinbami2000@yahoo.com

Received 06-V-2005. Corrected 15-VIII-2005. Accepted 30-IX-2005.

Abstract: A total of 360 randomly selected specimens of *Clarias gariepinus* (Clariidae) recovered from Lekki Lagoon were examined for intestinal helminth parasites. Parasite prevalence and worm burden were low; 17 (4.72%) of the specimens examined were infected with gastrointestinal helminths. The helminth worms recovered include, three cestodes *Polyonchobothrium clarias*, *Stocksia pujehuni* and *Wenyonia acuminata* and a nematode, *Paracamallanus cyathopharynx*. There was no statistically significant difference in the infection of the male and female *Clarias gariepinus*; the infection rates in male and female samples were 5.75% and 3.76% respectively. Parasite prevalence was related to the length and weight of the specimens. The fish samples were observed to show negative allometric growth and smaller samples recorded higher helminth infection. Rev. Biol. Trop. 54(1): 93-99. Epub 2006 Mar 31.

Keywords: Helminth parasites, *Clarias gariepinus*, Lagos.

Clarias gariepinus (Burchell, 1822) family Clariidae is generally considered to be one of the most important tropical catfish species for aquaculture in West Africa (Clay 1979). *Clarias* species inhabit calm freshwaters ranging from lakes, streams, rivers, swamps to flood plains, many of which are subject to seasonal drying. The most common habitats of the catfish are flood plains, swamps and pools. The catfish can survive during the dry seasons due to the possession of accessory air breathing organ (Bruton 1979a, Clay 1979). Since the last three decades, *Clarias gariepinus* has been considered to hold great promise for fish farming in Africa; the fish having a wide geographical spread, a high growth rate, resistant to handling and stress, and well appreciated in a wide number of African countries.

The males and females of *Clarias gariepinus* are readily distinguishable. The males possess a distinct sexual papilla that is

conspicuously located behind the anus. The sexual papillae are absent in females.

The fish is generally classified as omnivores or predators feeding mainly on aquatic insects, fish and higher plants debris as reported for catfishes in the River Ubangui, Central African Republic (Micha 1973). They have also been found to feed on terrestrial insects, molluscs and fruits. Bruton (1977b) found that adult catfishes in Lake Sibaya (South Africa) fed mainly on fish or crustacean, while terrestrial and aquatic insects were the important diet of juvenile and adult fishes inhabiting shallow areas. The catfishes utilize various kinds of food resources available in their habitats.

Documentation of parasite fauna of fishes especially used in aquaculture is on the increase. In northern Nigeria, a number of gastrointestinal helminth worms have been documented, in *C. gariepinus* by Oniye *et al.* (2004) in Zaria and Yakubu *et al.*

(2002) in Plateau state. Oniye *et al.* reported the occurrence of cestodes, *Anomotaenia* sp. (2.5%), *Monobothrium* sp. (13.33%) and *Polyonchobothrium clarias* (1.67%); the nematode, *Procamallanus laevionchus* and an acanthocephalan, *Neoechinorhynchus rutli* (0.83%).

As a consequence of the wide geographic spread, the diverse and non-specific food types or diet, the commercial and aquaculture values of the catfish, *C. gariepinus*, an investigation of the helminth fauna was undertaken to evaluate the nature and impact of helminth infections in the southern part of the country. The present study was carried out in Lekki Lagoon which serves as an important fishing zone to the inhabitants of Epe and metropolitan Lagos at large.

MATERIALS AND METHODS

Study Area: Lekki lagoon supports a major fishery in Nigeria. The Lekki lagoon located in Lagos State Nigeria lies between longitudes 4°00' and 4°15' E and between latitudes 6°25' and 6°37' N, has a surface area of about 247 km² with a maximum depth of 6.4 m. A large portion of the lagoon is shallow and less than 3.0 m deep. The Lekki lagoon is part of an intricate system of waterways made up of lagoons and creeks that are found along the coast of South-western Nigeria from the Dahomey border to the Niger Delta stretching over a distance of about 200 km. It is fed by the River Oni discharging to the North-eastern and the Rivers Oshun and Saga discharging into the North-western parts of the lagoon.

The vegetation around the lagoon is characterised by shrub and raphia palms, *Raphia sudanica*, and oil palms *Elaeis guineensis*. Floating grass occur on the periphery of the lagoon while coconut palms *Cocos nucifera* are widespread in the surrounding villages.

The lagoon which experiences both dry and rainy seasons typical of the southern part of Nigeria supports a major fishery in Nigeria. The rich fish fauna of the lagoon includes *Heterotis niloticus*, *Gymnarchus niloticus*, *Clarias gariepinus*, *Malapterurus electricus*,

Synodontis clarias, *Chrysichthys nigrodigitatus*, *Channa obscura*, *Mormyrus rume*, *Calabaricus calamoichthys*, *Tilapia zilli*, *Tilapia galilae*, *Hemichromis fasciatus* and *Sarotherodon melanotheron* (Kusemiju 1981). The map of Lekki Lagoon is shown in figure 1.




Fig. 1. Map of Lekki Lagoon.

Collection and Examination of specimens for Parasites: From early, 2003 to late, 2004, 360 randomly selected fresh specimens of *Clarias gariepinus* recovered from Lekki lagoon were purchased at Oluwo Market at Epe, Lagos, Nigeria and were examined for parasites. The weights, standard lengths and total lengths of the fishes were recorded. The fishes were dissected and the alimentary canals were removed and cut into parts in physiological saline for parasite recovery. The intestines were further carefully slit open to aid the emergence of parasites. The recognition of the worms was enhanced by the wriggling movements on emergence.

Processing of Parasites Recovered: The recovered helminth parasites were fixed in 70% alcohol, counted and recorded. Whole mount histological preparations of worms stained with Haematoxylin and eosin were

prepared. Identification of specimens to species level was undertaken and confirmed at the British Museum (Natural History), United Kingdom.

RESULTS

Three hundred and sixty specimens of *Clarias gariepinus* were subjected to parasitologic investigations. All helminthic infections observed and recorded were restricted to the intestine. Three gastrointestinal helminth worms recovered included a nematode, *Paracamallanus cyathopharynx* (Baylis, 1923) (Camallanidae) and three cestodes, *Polyonchobothrium clarias* (Woodland, 1925) (Ptychobothriidae), *Wenyonia acuminata* (Woodland, 1923) (Caryophyllaeiidae) and *Stocksia puehuni* (Woodland 1937) (Caryophyllaeiidae). No acanthocephalan was recovered.

Seventeen samples were infected. A prevalence of 4.72% was recorded. Single and mixed infections with 2 helminth worms were observed. Table 1 presents the prevalence of helminthic infections with the male showing a slightly higher prevalence (5.75%) than the females (3.76%). The variation in size of the catfishes in relation to helminthic infection of *Clarias gariepinus* is presented in Table 2. Fishes in length categories 16-20 cm and 26-30 cm recorded significantly higher helminth prevalence than the other length categories. The prevalence of parasitic infections was minimal in the length group 10-15 cm and 21-25 cm.

The weight of the fish samples examined ranged from 13.46 g to 160.33 g. Parasitic helminthes were observed in all weight categories. The results also showed that smaller specimens are more liable to infections than the bigger ones. Figures 2, 3, 4 and 5 showed scattered diagrams of total length against weight of males, females and combined sexes. The specimens of *Clarias gariepinus* showed negative allometric growth; this implies that growth is proportional to increase weight. Observations indicate that the smaller fishes recorded higher infection than bigger fishes.

DISCUSSION

The gastrointestinal helminth parasites of *Clarias gariepinus* in Lekki lagoon, Lagos, Nigeria were investigated. A prevalence of 4.72% gastrointestinal helminthic infections

TABLE 1
Prevalence of intestinal helminth infection in relation to sex of Clarias gariepinus in Lekki lagoon

	Male	Female	Combined sex
Number examined	174	186	360
Number infected with parasite	10	7	17
Percentage of infection	5.75	3.76	4.72

Chi-square value = 3.841

TABLE 2
Intestinal helminth infection in relation to size of Clarias gariepinus, in Lekki lagoon

	Group length				Total
	10-15 cm	16-20	21-25	26-30	
Number examined (%)	72 (20)	223 (61.9)	51 (14.17)	14 (3.89)	360 (100)
Number infected with parasite	3	10	2	2	17
Percentage of infection	4.16	4.48	3.92	14.29	4.72

Chi-square value = 7.814




Fig. 2. Scattered diagram total length against total weight of *Clarias gariepinus* (combined sexes).




Fig. 3. Log total length / Log weight of *Clarias gariepinus* from Lekki lagoon (Male).




Fig. 4. Log total length / Log weight of *Clarias gariepinus* from Lekki lagoon (Female).




Fig. 5. Log total length against Log weight of *Clarias gariepinus* from Lekki lagoon (combined sexes).

was recorded. The parasites recovered from the present study are *Polyonchobothrium clarias*, (Woodland 1925) *Stocksia pujehuni*, (Woodland 1937) *Paracamallanus cyathopharynx* (Baylis 1923) and *Wenyonia acuminata* (Woodland 1923). According to the host parasite checklist on African freshwater fishes of Khalil and Polling (1997), *P. clarias* has been documented in *Clarias gariepinus*. However, the present study is the first record of *P. cyathopharynx* and *Wenyonia acuminata* in *Clarias gariepinus*. Oniye *et al.* (2004) in Zaria, Nigeria, isolated five species of helminth parasites comprising three cestodes, one nematode, and one acanthocephala. In the present study no acanthocephalan species was recorded. The parasite fauna of *C. gariepinus* varied from the southern and northern hemisphere of the country. This is significant from the aquaculture viewpoint.

Diverse prevalence of *Polyonchobothrium clarias* are documented in the literature. Imam (1971), reported an incidence of *Polyonchobothrium clarias* as high as 42% for *Clarias gariepinus* dwelling in the Nile, in mid Egypt (Cairo and Guiza provinces) while Sawa (1982) recorded the prevalence of 22% in Manzalla Lake, Egypt. However, a low parasite prevalence and burden were observed in this

survey. Yakubu *et al.* (2002) in a comparative study of the helminths of *Tilapia zilli* (Gervais, 1848) and *Clarias gariepinus* from River Uke, Plateau State, Nigeria, reported a 55% prevalence of helminth infections in *Clarias gariepinus*. The nature of the lagoon, inclusive of its physical and physico-chemical parameters may influence the low prevalence of intestinal helminth in the catfishes compared to those of inland waters. Climatological factors which influence the fauna and flora of the water body and hence the feeding patterns and habits of the fishes may be responsible for the variable parasite fauna and intensity in the different zones in the country. While the climate in the south is equatorial the northern zone experiences savannah conditions. Geo-climatic factors apart from influencing the prevalence and intensity of the helminthic infection probably play a significant role in the parasite fauna of *Clarias gariepinus* in Nigeria.

Paracamallanus cyathopharynx is an ovoviviparous camallanid nematode whose larvae are liberated into the gut of the host and pass out with the faeces (Moravec 1974). The first moult of the parasite takes place in the copepod intermediate host and the last two moults take place in the fish. It is speculated that the utilization of the copepods as food

by the catfishes in the lagoon is minimal and higher in the small sized fishes; other food types being readily accessible for fish consumption. Available physical space, the geographical position, physico-chemical factors of the water body, the nature of diet will influence the nature and the frequency and intensity of helminth infections in the aquaculture of catfishes.

In the present study, *Wenyonia acuminata* was isolated from the intestine of *Clarias gariepinus*. In West Africa, species of *Wenyonia* have been reported in *Synodontis* species (Khalil and Polling 1997). For instance, Banhawy *et al.* (1975) isolated *Wenyonia virilis* from the ileum of the silurid fish, *Synodontis schall*.

The occurrence of these parasites also in *Clarias gariepinus* may be indicative of similar diets, feeding habits and patterns amongst the freshwater fishes. The documented host for *Wenyonia acuminata* include *Synodontis membranaceus*.

It is important to note that many helminth parasites in the alimentary canal of fish are pathogenic. The pathological effects of these helminths are as a result of the mechanical damage caused by the attachment organs. *Polyonchobothrium clarias* has been known to induce series of pathological lesions on the intestines of *Clarias gariepinus*. Banhawy *et al.* (1975) reported degenerative changes in the gut wall of *Synodontis schall*. Some pseudophylid cestodes are known to cause irritation of gastric mucosa of fish (William 1969, Euzet 1959). The impact of worm burden and prevalence on fish size needs to be assessed.

Parasites were recovered from fishes in all weight categories. The report of Oniye *et al.* (2004) indicated that fishes with weight (150-299.9 g) were free of infection. Also, the higher prevalence of helminthic infection in fishes weighing between 350.0-399.0 g and 450.0-500.0 g could be related to the larger sized fishes recovered in the north. Large-scale fishing in the Lekki Lagoon may account for the medium sized nature of the prevalent fishes compared to those from Zaria.

Variations in the infections of the different length categories were recorded. The prevalence of parasitic infections correlates with fish length which in turn corresponds to fish age as reported by Lagler *et al.* (1979). Low infection was found in fish length 10-15 cm (4.16%) and 21-25 (3.92%). This may be attributed to probable differences in feeding habits with fish age and immunity. Oniye *et al.* (2004) had also indicated the importance of changing feeding habits of the fish with age on the parasite prevalence and intensity.

Further studies are required to establish the diverse factors that influence the prevalence of gastrointestinal helminths, considering the consequent pathogenicity and the aquaculture value of *Clarias gariepinus* in Nigeria.

ACKNOWLEDGMENT

The authors express sincere gratitude to David Gibson of the British Museum, Parasitic Worms Division, United Kingdom, for identifying the parasites to species level.

REFERENCES

- Banhawy, M.A., M.F.A. Saoud, I.M. Anwar & M.K. El-Naffar. 1975. The histopathological effects of the parasitic tapeworm *Wenyonia virilis* on the ileum and liver of the silurid fish *Synodontis schall*. *Ann. Zool.* 11: 83-101.
- Bruton, M.N. 1979a. The breeding biology and early development of *Clarias gariepinus* (*Pisces claridae*) in Lake Sibaya, South Africa, with a review of breeding species of the subgenus *Clarias* (*clarices*). *Trans. Zool. Soc. London* 35: 1-45.
- Bruton, M.N. 1979b. The food and feeding behaviour of *Clarias gariepinus* (*pisces, claridae*) in Lake Sibaya, South Africa, with its emphasis on its role as a predator of cochleids. *Trans. Zool. Soc. London* 35: 47-114.
- Clay, D. 1979. Population biology, growth and feeding of the African Catfish, *Clarias gariepinus*, with special reference to juveniles and their importance in fish culture. *Arch. Hydrobiol.* 87(4): 453-482.

- Euzet, L. 1959. Recherches Sur les cestodes tetraphyllides des selaciens cotes de France. Thesis. University of Montpellier, 263 pp.
- Imam, E.A. 1971. Morphological and biological studies on the enteric helminths infecting some of the Egyptian Nile fishes particularly *Polyonchobothrium clarias* of *Karmot clarias lazera* and *Clarias anguillaris*. Ph. D. Thesis, Parasitology. Cairo University.
- Khalil, L.F. & L. Polling. 1997. Checklist of the helminth parasites of African freshwater fishes. University of the North Republic of South Africa, 161 pages.
- Kusemiju, K. 1981. The hydrobiology and fishes of the Lekki Lagoon, Nigeria. Nig. J. Nat. Sci. 3(1-2): 135-146.
- Lagler, K.F., J.E. Bardach & R.R. Miller. 1997. Ichthyology. John Wiley, New York.
- Micha, J.C. 1973. Etude des populations piscicoles de l'ubangui et tentative de selection et d'adaptation de quelques especes a l'etang de pisciculture. Centre Technique Forestiere Tropical, Nogent Sur Marne, 100 pp.
- Moravec, F. 1974. The development of *Paracamallanus cyathopharynx* (Baylis, 1923) Nematoda: Camallanidae. Folia Parasit. 21: 333-343.
- Oniye, S.J., D.A. Adebote & O.I. Ayanda. 2004. Helminth parasites of *Clarias gariepinus* in Zaria, Nigeria. J. Aquatic Sci. 19(2): 71-76.
- Sahlab, A. 1982. Studies on the enteric helminth parasites of fish from Lake Manzalla, M.Sc. Thesis (Parasitology). Cairo University, Egypt.
- Yakubu D.P., E. Omoregie & J.W. Wade. 2002. A comparative study of gut helminths of *Tilapia Zilli* and *Clarias gariepinus* from river uke, plateau state, Nigeria. J. Aquatic Sci. 17(2): 137-139.
- Williams, H.H. 1969. Some observations on *Parabothrium gadipollachii* (Rudolphi, 1810) and *Abothrium gadi* van Beneden, 1870 (Cestoda: Pseudophyllidae) including an account of their mode of attachment and variation in the two species. Parasitology: 303-322.

