

Adverse reactions from community directed treatment with ivermectin (CDTI) for onchocerciasis and loiasis in Ondo State, Nigeria

O.A.Otubanjo¹, G.O.Adeoye¹, C.A.Ibidapo², B.Akinsanya¹, P.Okeke¹, T. Atalabi¹, E.T. Adejai³ & E. Braide⁴

1. Department of Zoology, University of Lagos, Akoka, Yaba, Lagos, Nigeria; adetoro2001@yahoo.com, goa4567@yahoo.com, akinbami2000@yahoo.com
2. Department of Zoology, Faculty of Science, Lagos State University, Ojo, Lagos, Nigeria; jokeibidapo@yahoo.com
3. Onchocerciasis Section, Ministry of Health, Ondo State, Nigeria.
4. Department of Biological Sciences, University of Calabar, Calabar, Cross River State, Nigeria; ekanem_b@hotmail.com

Received 07-VI-2007. Corrected 30-VI-2008. Accepted 31-VII-2008.

Abstract: Onchocerciasis is an endemic disease in Ondo state, Nigeria. Community directed distribution of ivermectin is currently on-going in some local government areas of the state. Randomly selected persons (2 331 males and 2 469 females) were interviewed using a modified rapid assessment procedure for *Loa loa* (RAPLOA) to assess community directed treatment with ivermectin. The retrospective study evaluated the coverage, impacts and adverse reactions to the drug treatment. A questionnaire was administered by house-to-house visit in six local government areas, implementing community directed treatment with ivermectin (CDTI) in this bioclimatic zone. A total of 2,398 respondents were reported to have participated in the treatment. The overall ivermectin coverage of 49.96% was recorded (range 0 – 52% in different communities). Adverse reactions from ivermectin administration were experienced in 38% of individuals. Diverse adverse reactions experienced included predominantly itching (18.50%); oedema, especially of the face and the limbs (8.2%); rashes (3.4%) and body weakness (2.4%). Expulsion of intestinal worms occurred in 0.96% of the respondents. The occurrence of adverse reactions in relation to age categories was statistically significant. Neither fatal nor severe adverse reactions were reported by respondents. Significantly, despite experienced adverse reactions, continued participation, acceptability and compliance to ivermectin treatment was expressed by the various communities. This attitude is in consonance with the African Programme for Onchocerciasis Control (APOC) objectives. Rev. Biol. Trop. 56 (4): 1635-1643. Epub 2008 December 12.

Key words: onchocerciasis, loiasis, adverse reactions, community directed treatment with ivermectin, rapid assessment procedure for *Loa loa*, African Programme for Onchocerciasis Control.

Onchocerciasis or river blindness, a major public health problem, afflicts 18 million people world wide, of whom 99% live in Africa south of the Sahara (Ety'ale 2001, 2002). The control of onchocerciasis is under the WHO programme of the African Programme for Onchocerciasis Control (APOC). This programme entails mass chemotherapy by community directed treatment with ivermectin (CDTI), following the free donation of the drug (Mectizan ®), by the manufacturer, Merck &

Co. Inc. The use of ivermectin revolutionized onchocerciasis control in Africa and the control programme is geared to reduce the transmission of the disease.

Meanwhile, a single annual dose treatment with ivermectin by community directed distributors forms the principal intervention in the control of the disease, Mild adverse reactions, including pruritus (itching), fever and rashes within the first 2-3 days of ivermectin administration have been variously documented

(Gardon *et al.* 1997, Kipp 2003, WHO 2003). However, the diversity of adverse symptoms is complicated in regions of co-endemicity of onchocerciasis with loiasis. Serious complications, such as severe and sometimes fatal encephalopathic adverse reactions and coma in patients with onchocerciasis combined with high intensity of *Loa loa* have been reported (Chippaux *et al.* 1996, Gardon *et al.* 1997, Boussinesq *et al.* 1998). Also, ivermectin is contra-indicated in patients with trypanosomiasis and central nervous disorders, especially meningitis. In the absence of a clinical macrofilaricide, the risk of severe reactions threatens the success of the control programme in certain African communities with endemic loiasis.

Using rapid epidemiological mapping of onchocerciasis, the pattern of onchocerciasis endemicity in Ondo State, South western Nigeria was undertaken in 1994 by the State task force. Ondo State is in a bioclimatic zone, ranging from rainforest to savannah mosaic. It is highly endemic for onchocerciasis. Ivermectin administration started in 1994 under the auspices of the United Nations Children Education Fund in five local government areas. In June 2000, the African Programme for Onchocerciasis Control and communities directed treatment with ivermectin commenced in sixteen onchocerciasis endemic local government areas in the State.

MATERIALS AND METHODS

Study site: The study involved 60 communities in six selected local government areas in Ondo State, South Western Nigeria. The studied local government areas were Owo, Akure North, Ifedore, Akure South, Ondo East and Ondo West. The projected population of the state is about 2.9 million. Ondo state covers an area of 15 600 square kilometers, lying in the bioclimatic zone, ranging from rainforest to forest, savanna mosaic to guinea savannah and mountainous areas. The study areas were randomly selected within the onchocerciasis endemic areas under mass ivermectin treatment. The main occupations in the State are farming and trading. Advocacy was established

at all levels of government, prior to the commencement of the study.

A total of 4 800 individuals (2 331 males and 2 469 females), 18 years old and above were randomly selected and interviewed during the survey. A modified rapid assessment procedure for loiasis (TDR 2000) was administered by house to house visits in the study areas. The rapid assessment procedure for loiasis was used in establishing the local name, the history and occurrence of loiasis in the studied communities. The year/years of treatment and impacts of ivermectin administration were recorded for individuals interviewed.

Statistical analysis: The data obtained were statistically analyzed using the Chi-square test.

RESULTS

Of the 4 800 individuals, 2 398 were reported to have participated in the community directed treatment with ivermectin between the periods 1996 to 2004 (Table 1). Amongst these participants, ivermectin had been administered only once in 1 771 (73.84 %) of the treated respondents. The overall coverage of 49.96%, ranging from 0 – 52% was reported in the different communities (Table 2). Participation and coverage in the community directed treatment with ivermectin was highest in Ifedore Local Government Area and least in Owo Local Government Area (Table 3). Both genders participated in a 1: 1 ratio in the community directed treatment with ivermectin within the various local government areas (Table 3). This study revealed that the age groups 21 – 30 and 31 – 40 were actively involved in community directed treatment with ivermectin programme in the State (Table 2). It was noted that participation in the community directed treatment with ivermectin was significantly age-related in some local government areas.

Adverse reactions to ivermectin experienced within the communities ranged from pruritus (itching); swelling or oedema of the body, particularly the face, stinging, general body pain, muscular or joint pain, rashes, body stiffness and general malaise occurring within

TABLE 1
Summary of CDTI coverage from 1989-2004 in the studied population

Duration	Female (%)	Male (%)	Total
1989	-	1 (100.00)	1
1996	-	1 (100.00)	1
1997	2 (33.33)	4 (66.67)	6
1998	10 (35.71)	18(64.29)	28
1999	21 (39.62)	32 (60.38)	53
2000	43 (42.57)	58 (57.43)	101
2001	130 (49.81)	131 (50.19)	261
2002	302 (47.86)	329 (52.14)	631
2003	702 (54.00)	598 (46.00)	1300
2004	11 (68.75)	5 (31.25)	16
Total	1221 (50.92)	1177 (49.08)	2398
WithoutCDTI	1248 (51.96)	1154 (48.04)	2402
Overall Total	2469 (51.44)	2331 (48.56)	4800

the first week of drug administration. These reactions were indicated to extend in some respondents for a period of 7-14 days. Adverse reactions occurred in various combinations. In most cases, 914 (38.15%) respondents' experienced adverse reactions on ivermectin administration (Table 4).

The predominant overall adverse reaction in each community was itching (18.5%). Itching occurring for 1-5 days was usually accompanied by other symptoms and may extend for two weeks in some areas. Swelling or oedema of the body, particularly the face (8.1%), rashes (3.4%), boils (3.4%), headache and fever (0.72%) were experienced by respondents in the various communities (Table 5). Generally, adverse reactions were experienced within 1-7 days. Stooling was experienced by 0.40% of individuals after ivermectin administration. The degree of adverse reactions varied in the different local government areas (Table 6), with Ifedore Local Government Area recording the highest level of occurrence of adverse reactions. Ondo East Local Government Area had the lowest record of adverse reactions.

Adverse reactions were high in the 21 – 30 years age group with 154 (22.4%) individuals. This was statistically significant ($P>0.05$). The age groups 50- 70 years and above recorded significant adverse reactions. However, adverse reactions were not significant in relation to the gender of the respondents. Within the gender, adverse reactions occurred in 51.10% and 48.80% males and females respectively. The ratio of male to female with adverse reactions was 1:1.06. Consequent to ivermectin treatment some respondents (0.96%) experienced intestinal worm expulsion.

Loiasis endemicity in the communities ranged from low to moderate. The common name for loiasis in all the communities is “*aran oju*” (meaning eye worm).

DISCUSSION

Community directed treatment with ivermectin (CDTI) is the principal drug delivery strategy for onchocerciasis control. In Ondo State, ivermectin treatment control programme commenced in 1994, under the auspices of

TABLE 2
CDTI coverage by age group in the six LGAs in Ondo state

AGE GROUP (Yrs)	AGE GROUP (Yrs)						TOTAL		
	15 -20	21 -30	31 -40	41 - 51	51 - 60	61 -70		>71	
Owo (01)	Examined	151	223	138	109	74	59	25	800
	No (%)	46 (30.46%)	57 (25.56%)	51 (36.96%)	42 (38.53%)	33 (44.59%)	25 (42.37%)	15 (60.00%)	269 (33.63%)
Akure North (02)	Examined	112	234	164	99	89	59	23	800
	No (%)	21 (18.75%)	43 (18.38%)	40 (24.39%)	33 (35.35%)	37 (41.57%)	27 (45.76%)	8 (34.78%)	211 (26.38%)
Ifedore (03)	Examined	177	239	112	112	87	41	24	800
	No (%)	75 (42.37%)	105 (43.93%)	74 (66.07%)	89 (79.46%)	67 (77.01%)	25 (60.98%)	14 (58.33%)	449 (56.13%)
Akure South (04)	Examined	132	212	175	102	71	44	26	800
	No (%)	40 (30.30%)	73 (34.43%)	74 (42.29%)	55 (53.92%)	47 (66.20%)	23 (52.27%)	15 (57.69%)	327 (40.88%)
Ondo East (05)	Examined	204	246	142	87	61	48	24	800
	No (%)	49 (24.02%)	49 (19.92%)	33 (23.24%)	23 (26.44%)	12 (19.67%)	18 (37.50%)	8 (33.33%)	192 (24%)
Ondo West (06)	Examined	156	252	144	89	73	60	25	800
	No (%)	37 (23.72%)	83 (32.94%)	65 (45.14%)	48 (53.93%)	36 (49.32%)	35 (58.33%)	13 (52.00%)	317 (39.63%)
Total Examined No. Positive (%)	Total Examined	932	1406	875	598	455	311	147	4800
	No. Positive (%)	268 (28.76%)	410 (29.16%)	337 (38.51%)	292 (48.83%)	232 (50.99%)	153 (49.20%)	73 (49.66%)	1765 (36.77%)

TABLE 3
CDTI coverage by sex in the six local government areas in Ondo State

		CDTI COVERAGE RATES		
		Male	Female	TOTAL
Owo (01)	Examined	345	455	800
	No. (%) Positive	161 (45.87)	190 (54.13)	351 (43.88)
Akure North (02)	Examined	443	357	800
	No. (%) Positive	271 (61.04)	173 (38.96)	444 (55.50)
Ifedore (03)	Examined	343	457	800
	No. (%) Positive	93 (27.10)	120 (26.30)	213 (26.63)
Akure South (04)	Examined	409	391	800
	No. (%) Positive	219 (53.50)	187 (47.80)	406 (50.75)
Ondo East (05)	Examined	395	405	800
	No. (%) Positive	128 (32.40)	143 (35.31)	271 (33.87)
Ondo West (06)	Examined	396	404	800
	No. (%) Positive	177 (44.70)	174 (43.10)	351 (43.88)
TOTAL	Examined	2469	2331	4800
	No. (%) Positive	1177 (47.80)	1221 (52.40)	2398 (49.95)

TABLE 4
Percentage adverse reactions in the six Local Government Areas in Ondo State

Local government areas	CDTI Count	Adverse reactions Count	% Adverse reactions Count
Owo LGA.	351	128	36.47
Akure North LGA	444	257	57.88
Ifedore LGA	587	224	38.16
Akure South LGA	394	106	26.90
Ondo East LGA	271	79	29.15
Ondo West LGA	351	120	34.19
TOTAL	2398	914	38.12

United Nations in five local government areas. In June 2000, the African Programme for Onchocerciasis Control and ivermectin strategy became operational. The treatment coverage is still low. This report affirms a 49.95% community directed treatment with ivermectin in the study areas. African Programme for Onchocerciasis Control recommends at least

a target of 65% therapeutic coverage annually. This study reveals that 2 402 (50.04%) of the respondents have not received ivermectin in the study area. A small percentage of the communities had achieved the target therapeutic coverage. A major reason for this is that community directed treatment with ivermectin is irregular and haphazard.

TABLE 5
Adverse reactions experienced within the studied population

Adverse Reaction	Local Government Areas in %						Total %
	1	2	3	4	5	6	
Itching	16.43	3.83	21.80	13.71	10.70	17.38	17.91
Swelling	6.52	2.92	3.92	6.60	9.59	10.54	7.93
Stooling	0.57	-	0.34	0.51	0.74	-	0.40
Body Pain	1.70	0.23	0.85	1.02	-	1.14	0.85
Headache and Fever	-	0.23	0.85	0.51	0.74	-	0.72
Weakness	1.42	0.90	-	3.05	2.95	1.99	2.29
Boils/Rashes	3.97	0.90	3.07	2.03	2.95	5.41	3.30
Body Stiffness	0.85	-	0.17	-	-	-	0.28

TABLE 6
Adverse reactions in the different Local Government Areas

LOCAL GOVERNMENT	COUNT	Adverse Reactions YES	% WITHIN LG CODE	% WITHIN Adverse Reactions
OWO	800	100	12.50%	7.75%
AKURE NORTH LGA	800	393	49.13%	30.47%
IFEDORE LGA	800	529	66.13%	41.00%
AKURE SOUTH	800	90	11.25%	6.98%
ONDO EAST	800	60	7.50%	4.65%
ONDO WEST	800	118	14.75%	9.15%
TOTAL	4800	1290(53.79%)		

N.B 2,398 respondents participated in CDTI.

Several problems currently associated with drug distribution coverage in the state includes, instability and disruption of the programme by state bureaucratic processes, fuel scarcity, fund disbursement and general lack of adequate commitment to the programme by local government coordinators. The lack of incentives for the community directed distributors and inaccessibility of some target communities are additional obstacles to drug distribution.

The success of African Programme for Onchocerciasis Control will depend on the sustenance of community directed treatment with ivermectin (Amazigo *et al.* 2002)

Adverse reactions to ivermectin occurred in 53.79% of respondents that participated in the drug treatment. There is no gender difference in their participation in the community directed treatment with ivermectin. Gender issues relating to community directed treatment with ivermectin constitute a challenge to African Programme for Onchocerciasis Control (Clemmons *et al.* 2002, Seketeli 2002). Within the study period, 1 221(50.92%), females participated in the programme. Many respondents, 1 771 (73.85%) had only been administered the drug once, since the inception of the programme. This report establishes the low level

and haphazard distribution of ivermectin in the study areas. However, the African Programme for Onchocerciasis Control recommends several years of annual ivermectin administration to effectively establish reduced transmission of onchocerciasis and thereby intercept the disease endemicity in the African region.

Adverse reactions experienced are varied and similar to those that has been documented in previous studies (Zea-Flores *et al.* 1992, Burnham 1993, Baraka *et al.* 1995, Kipp *et al.* 2003). Significantly itching, oedema, boils and rashes were the reported adverse reactions. Host inflammatory responses had been indicated due to dying microfilariae and not by direct drug toxicity (Turner *et al.* 1994). No fatal, severe adverse reaction was reported in all the communities. The attitudes of individuals to adverse reactions and further treatment were non-conflicting. Similar reports have been documented (Baraka *et al.* 1995). Rapid procedure for loiasis is a useful tool in the continued rapid assessment of community prevalence of loiasis infection especially in areas of co-infection of loiasis with onchocerciasis.

According to the data, the prevalence of loiasis in all the communities was much below 40%. The low endemicity of loiasis is considered to have low risk of adverse reactions during mass drug therapy with ivermectin (Tropical Disease Research 2002). The level of loiasis endemicity varied considerably between the local government areas and communities. The bioclimatic conditions of the different local government areas will affect the vector habitat, thereby impacting on the occurrence of the disease. While every community had a local name for the eye worm, Calabar swelling “*awoka*” (meaning moving about) was not common and tended to be less specific in the study areas.

Ondo State is hypoendemic for loiasis (Ibidapo *et al.* pers comm) and thereby ivermectin administration is considered relatively safe, despite some degree of reported adverse reactions. Severe or fatal adverse reactions were non-existent.

The need to monitor adverse reactions following repeated ivermectin treatment is

important to achieve the goals and objectives of African Programme on Onchocerciasis Control. Reduced adverse reactions from first to sixth rounds of treatments (40.6% to 15.6%) in 890 individuals in Kwara State, in Nigeria have been recorded (Oyibo and Fagbenro-Beyioku 2003). They concluded that adverse reaction rates did not affect future participation in community directed treatment with ivermectin, as adequate community mobilization with health education messages were in place. Similar reports were documented by Kipp *et al.* 2003 and Baraka *et al.* 1995. The pretreatment density of microfilariae in the skin influences the occurrence and intensity of adverse reactions. Continued drug delivery has been indicated to result in diminishing adverse reactions over time (Zea-Flores *et al.* 1992, Burnham 1993). Despite some degree of adverse reactions experienced by respondents in this study, participation, acceptability and compliance to ivermectin administration was reasonable in consonance with the programmes’ objectives.

The efficacy and action of ivermectin against filarial infections especially lymphatic filariasis, many intestinal parasites, lice and scabies have been documented. Zea – Flores *et al.* 1992 reported 38% worm expulsion on ivermectin administration in Guatemela. While this is considered to be an added advantage in ivermectin administration; affected individuals in many of the communities considered worm expulsion to be an adverse reaction. It is believed amongst the rural communities that a ‘healthy’ individual requires some degree of intestinal worm infection. Socio-culturally, worm expulsion was considered to be an adverse reaction by some rural respondents.

The occurrence of adverse reactions was predominant in the age group 21 -30. This may probably be due to the degree of exposure of ivermectin to this age bracket or may be influenced by the level of parasitism, since the intensity of microfilariae in the skin influence the nature and severity of adverse reactions.

This study revealed the need to increase ivermectin distribution to effectively achieve the targeted goals of the programme. The low

endemicity of *Loa loa* in the study area permits the administration of ivermectin. A biannual mass drug administration in Ondo state has been recommended by Idowu 2004 as a consequence of the high endemicity of the disease in the region.

ACKNOWLEDGMENTS

This investigation was financially supported by Sight Savers International, Nigeria. We are grateful to the Onchocerciasis State coordinator, for his personal commitment to the success of this project.

RESUMEN

La oncocercosis es endémica en el estado Ondo, Nigeria. Se seleccionaron 4 800 personas al azar para evaluar con encuesta retrospectiva la cobertura, efectos y reacciones al tratamiento farmacológico con ivermectina administrado por la misma comunidad. La cobertura global de ivermectina fue 50 % con reacciones adversas en 38 % de los individuos. Estas fueron comezón picazón (18%), edema, especialmente de la cara y las extremidades (8%), erupciones cutáneas (3%) y debilidad (2%); dependieron de la edad y no hubo reacciones más graves. La expulsión de las lombrices intestinales se produjo en 96% de los encuestados. A pesar de las reacciones adversas, hubo continuidad, aceptación y cumplimiento del tratamiento con ivermectina, en consonancia con los objetivos del Programa Africano para el Control de Oncocercosis (APOC).

Palabras clave: oncocercosis, loiasis, reacciones adversas, ivermectina, procedimiento de evaluación rápida para *Loa loa*, Programa Africano para el Control de Oncocercosis.

REFERENCES

Amazigo, U.V., O.M. Obono, K.Y.Dadzie, J.Remme, J.Jiya, R.Ndyomugenyi, I.B.Rongon, M.Noma & A.Seketeli. 2002. Monitoring community directed treatment programmes for sustainability: lessons from the African Programme for Onchocerciasis Control (APOC). *Ann Trop Med Parasit.* 96: 575 – 592

Awadzi K., N.O.Opokn, E.T.Addy & B.T.Quartey. 1995. The chemotherapy of onchocerciasis XIX: The chemical and laboratory tolerance of high dose ivermectin. *Trop Med Parasitol.* 46: 31 - 7

Burnham, G.M. 1993. Adverse reactions to ivermectin treatment for onchocerciasis: results of a placebo

–controlled, double –blind trial in Malawi. *Trans R Soc Trop Med Hyg.* 87: 313 – 7

Baraka O.Z., A.K.Khier, K.M. Ahmend, M.M. Ali., A.E. el Mardi, B.M. Mahmoud, M.H.Ali., M.M.Homeida & J.F.Williams.1995. Community based distribution of ivermectin in eastern Sudan: acceptability and early post-treatment reactions. *Trans R Soc Trop Med Hyg.* 9: 316-8.

Boussinesq M., I. Gardon , N.Gardon-Wendel, I.Kamgno, P.Ngoumou & I.P.Chippaux 1998. Three probable cases of *Loa loa* encephalopathy following ivermectin treatment for onchocerciasis. *Am J Trop Med Hyg.* 58: 461-469.

Boussinesq M., I.Gardon., I.Kamgno & S.D.S. Pion. 2001. Relationship between the prevalence and intensity of *Loa loa* infection in the Central Province of Cameroon. *Ann Trop Med Parasit.* 95: 495-507.

Clemmons L., U.V.Amazigo, A.C.Bissek, M.Noma, U.Oyene, U.Ekpo, J.Msuya-Mpanju., S.Katenga & A. Seketeli. 2002. Gender issues CDTI of the African Programme for Onchocerciasis Control (APOC). *Ann Trop Med Parasit.* 96: S59-S74.

Chippaux J.P., M.Boussinesq, J. Gardon, N.Gardon-Wendel & J.C.Ernould .1996. Severe adverse reaction risk during mass treatment with ivermectin in loiasis-endemic areas. *Parasitol. Today.*12: 448-450.

Clemmons L, U.V.Amazigo, A.C. Bissek, M.Noma, U.Oyene, J.Msuya-Mpanju, S.Katenga & A.Seketeli. 2002. Gender issues in community –directed treatment with ivermectin CDTI of the African Programme for Onchocerciasis Control (APOC). *Ann Trop Med Parasit.* 96: S59-S74.

Ety’ale D. 2001.Vision 2020: update on onchocerciasis community. *Eye Health.* 14: 19 – 20.

Ety’ale D. 2002. Eliminating onchocerciasis as a public health problem: the beginning of the end. *Brit J Opthamol.* 86: 844 – 846

Gardon J., N.Gardon-Wendel, Demangangangue, J.Kamgno, J.P.Chippaux & M. Boussinesq. 1997. Serious reactions after mass treatment of onchocerciasis with ivermectin in an endemic area for *Loa loa* infection. *Lancet.* 350: 18-22.

Idowu E.T. 2004. Epidemiological, clinco-parasitological and control studies of onchocerciasis in selected communities of Ondo state, Nigeria. Ph.D. Thesis, University of Lagos, Nigeria.

KippW., J.Bamhuhiiga, T.Rubaale & D.W.Buttner. 2003. Adverse reaction to ivermectin treatment in *Simulium neavei*-transmitted onchocerciasis *Am J Trop Med Hyg.*69: 621-3.

- Oyibo W.A. & A.F. Fagbenro-Beyioku. 2003. Adverse reaction following annual ivermectin treatment of onchocerciasis in Nigeria. *Int J Infect Dis*. 7: 156 -9
- Seketeli A., G. Adeoye, A.Eyamba, E.Noruka , P.Drameh, U.V. Amazigo, M.Noma, F.Agboton, Y.Ahoton , O.Kale & K.Y. Dadziek. 2002 The achievement and challenges of the African programme for onchocerciasis control (APOC). *Ann Trop Med Parasit*. 96: S15-S28.
- Turner,P.F., K.A.Rocket, E.A.Otlesan.,H.Francis, K.Awadzi & I.Clark. 1994. Inter leukin-6 and tumor necrosis factor in the pathogenesis of adverse reactions after treatment of lymphatic filariasis and onchocerciasis. *J Infect Dis* 69: 1071 – 5.
- Tropical Disease Research 2000. World Health Organization (WHO) Implementation and sustainability of community directed treatment with ivermectin. Geneva.
- UNDP/ World Bank / WHO Special Programme for Research and Training in Tropical Disease. TDR / AFR / RP/96.1.
- Tropical Disease Research (TDR). 2002. Guidelines for rapid assessment of *Loa loa*. UNDP/World Bank/WHO.TDR/IDE/RAPLOA/02. 138 p.
- World Health Organization 2003. Report of a multicountry study: The involvement of community-directed distributors of ivermectin in other health and development activities. UNDP/World Bank/WHO TDR/IDE/CDD1/03.1.
- Zea-Flores R, Richards FO Jr. Gonzalez – Peralta C, Castro Ramirez J, Zea-Flores G., Collins RC, Cupp E.1992. Adverse reactions after community treatment onchocerciasis with ivermectin in Guatemala. *Trans R Soc Trop Med Hyg*. 86: 663-6.