

## GOBIESOCIDAE chupapiedras

*Tomicodon fasciatus*, chupapiedra de barras – Sin papilas en la región C del disco; coloración pardo claro con marcas pardo oscuro en el dorso y costados, extendiéndose a la región ventral (Guitart 1986, Briggs 1955).

*Acyrtops beryllinus*, chupapiedra verde – Un parche de papilas en forma de pera en la región C del disco; coloración verde esmeralda brillante con manchas y puntos blancos (Guitart 1986).

*Rimicola brevis*, chupapiedra enana – Disco pequeño, un parche de papilas en forma de barras en la región C del disco; coloración uniforme sin marcas oscuras (Briggs 1969).

*Gobiesox punctulatus*, chupapiedra punteada – Papilas aplanadas en la región C del disco en dos grupos bien separados; cabeza cubierta con manchas pardo oscuro, 2 barras blancas transversales que cruzan el dorso (Guitart 1986).

*Acyrtops rubiginosus*, chupapiedra roja – Papilas aplanadas en la región C del disco en dos grupos casi juntos; coloración roja uniforme o con bandas pardo oscuro transversales en la cabeza y cuerpo (Guitart 1986).

## BREGMACEROTIDAE bacaletes

*Bregmaceros atlanticus*, bacaleta antena – Aletas anal y segunda dorsal bilobuladas con un margen muy cóncavo; un solo radio filamentosos constituye la primera aleta dorsal (Goode & Bean 1895).

Apuntes / Notes:

## GOBIESOCIDAE clingfishes

*Tomicodon fasciatus*, barred clingfish – No papillae in disc region C; coloration light brown with dark brown markings on back, sides, extending to ventral surface.

*Acyrtops beryllinus*, emerald clingfish – A pear-shaped patch of papillae in disc region C; coloration bright emerald green with white spots and dots.

*Rimicola brevis*, pygmy clingfish – Disc small, a bar-shaped patch of papillae in disc region C; coloration uniform without dark markings.

*Gobiesox punctulatus*, stippled clingfish – Flattened papillae in disc region C in 2 widely separated groups; head covered with dark brown dots, 2 transverse white bars crossing the dorsum.

*Acyrtops rubiginosus*, red clingfish – Flattened papillae in disc region C in 2 closely approximated groups; coloration uniform red or with dark brown transverse bands crossing head and body.


## BREGMACEROTIDAE codlets

*Bregmaceros atlanticus*, antenna codlet – Anal and second dorsal fins bilobed with a deep concave margin; first dorsal fin a single filamentous ray.


## GOBIESOCIDAE




*Tomiodon fasciatus*




*Acyrtops beryllinus*



*Rimicola brevis*




*Gobiesox punctulatus*



*Acyrtus rubiginosus*



## BREGMACEROTIDAE



*Bregmaceros atlanticus*