

## Morphological and micromorphological characteristics of *Desmodium* fruits (Leguminosae: Papilionoideae)

Daiane M. Freitas<sup>1</sup>, Ademir Reis<sup>2</sup>, Roseli L. da Costa Bortoluzzi<sup>3</sup> & Marisa Santos<sup>4</sup>

1. Departamento de Botânica, Centro de Ciências Biológicas, Universidade Federal de Santa Catarina (UFSC), Campus Universitário, Florianópolis CEP 88040-900, Santa Catarina, Brasil and Herbário Barbosa Rodrigues (HBR), Av. Coronel Marcos Konder 800, Itajaí CEP 88301-302, Santa Catarina, Brasil; martinsf.daiane@gmail.com
2. Herbário Barbosa Rodrigues (HBR), Av. Coronel Marcos Konder 800, Itajaí CEP 88301-302, Santa Catarina, Brasil; ademir.reis.ufsc@gmail.com
3. Departamento de Engenharia Florestal, Centro de Ciências Agroveterinárias, Universidade do Estado de Santa Catarina (CAV-UDESC), Lages CEP 88520-000, Santa Catarina, Brasil; rosebortoluzzi@gmail.com
4. Departamento de Botânica, Centro de Ciências Biológicas, Universidade Federal de Santa Catarina (UFSC), Campus Universitário, Florianópolis CEP 88040-900, Santa Catarina, Brasil; marisa.santos@ufsc.br

Received 16-I-2014. Corrected 12-VII-2014. Accepted 14-VIII-2014.

**Abstract:** The genus *Desmodium* is represented in Santa Catarina State, Brazil, by 13 species, all with lomentaceous fruits. Shape, size and isthmus margin of loment vary, while the surface is glabrous, or covered by trichomes of different types. Morphological diversity of trichomes becomes particularly relevant to taxonomic description. The trichome types present on the surface of *Desmodium* fruits provide data for the identification and classification of species in the State. To assess this, three fruits of each species were collected and deposited at two herbaria, HBR and FLOR, in Santa Catarina, Brazil. Some rehydrated samples were examined using light microscopy (LM); and some sections were exposed to the following histochemical reagents: Sudan III for oils and Thionine for mucilage. The structural aspects of trichomes can be classified into uni- or multicellular and may still be simple, i.e., nonglandular or glandular. Using scanning electron microscopy (SEM), five types of trichomes were identified and analyzed among the *Desmodium* species studied: uncinata, uniseriate, globose multicellular, globose unicellular and subulate. Characteristics, such as loment margin and article form, glabrescent or pillous indument, trichome type, with or without papillous epidermal cells and epicuticular striations, showed relevant diagnostic value. An identification key was developed for *Desmodium* species from Santa Catarina State, Brazil, based on macro and micromorphological characters of the fruit. Rev. Biol. Trop. 62 (4): 1597-1608. Epub 2014 December 01.

**Key words:** dispersion, identification key, scanning electron microscopy (SEM), secretory structures, taxonomy, trichomes.

The *Desmodium* Desv. genus (Leguminosae, subfamily Papilionoideae; Ohashi, 2014) is included in the tribe Desmodieae and comprises about 524-530 species, with greater diversity in Southeast Asia, Mexico and South America (Ohashi, 2014). According to Lima, Oliveira and Tozzi (2014), 33 species are distributed throughout Brazil and 16 species in Santa Catarina State. The genus is characterized by shrubby or subshrubby habit, uni- or trifoliate leaves, lomentaceous or craspedium-type fruit

(whose pericarp splits into monospermous articles), and uncinata (hooked) trichomes, usually present on vegetative parts and fruits (Oliveira, 1983; Setubal, Lima, & Grings, 2010; Lima, 2011). In these species, the fruits can be sessile or stipitate with central isthmus, either eccentric or marginal, and the upper margin of the fruit, either straight or sinuous, but the lower margin is always sinuous.

According to Azevedo (1981), *Desmodium* species can have glabrescent or densely

pubescent fruit, and the trichomes can be uncininate or straight, uniseriate or multicellular, and long or short. Oliveira (1983, 1990) also cited different trichome types on the fruit surface for this genus: uncininate, straight glandular and straight nonglandular. Werker (2000) pointed out that trichomes are uni- or multicellular appendices originating from single epidermal cells present in various plant organs, such as vegetative parts, leaves, bracts, and roots near the apical meristem, or reproductive organs, including sepals, petals, stamens, gynoecium, fruit and seeds.

Epidermal surface ornamentation, such as conformations of the outer cell wall, trichomes and epicuticular waxes, contribute to maintaining inner water balance (Fahn, & Cutler, 1992). Convexity of epidermal cells in periclinal walls confers a greater ability to reflect solar radiation, with the walls acting as filters for the inner tissues, allowing adaptation to more xeric environments (Larcher, 2000). The amount of epicuticular waxes has been correlated to sunlight intensity (Barthlott, 1990); therefore, this type of deposition has been regarded as a taxonomic character (Barthlott et al., 1998). Since trichome density can change with environmental conditions, trichome type may be employed to identify a particular taxon.

Several studies have demonstrated the use of scanning electron microscopy (SEM) in angiosperm taxonomy, and many analyses have been performed on the surface of fruits and seeds in different plant groups. For example, Ávalos and Salinas (2003) found that trichomes of *Quercus* L. (Fagaceae) leaf surfaces could be used for delimiting species in that trichomes showed variation in shape, intensity, size and quantity of cells, as well as function, being either glandular or nonglandular. Agbagwa and Okoli (2005) used fruit epidermis micromorphology to study the systematics of *Abrus* Adans. (Papilionoideae). Ritter and Miotto (2006) used SEM to study fruit and seed surfaces for the taxonomy of 15 taxa of *Mikania* Willd. (Asteraceae). Akcin (2008) identified different microstructural patterns on the surface of seeds and fruits from

*Cynoglossum* L. (Boraginaceae) species. Kaya, Ünal, Özgökçe, Doğan, and Martin (2011) also used SEM to study fruit and seed surface characteristics to study the taxonomy of 11 genera of Brassicaceae.

The present research aimed to contribute to the identification and classification of the *Desmodium* genus from Santa Catarina State, Brazil, based on fruit macro- and micromorphology, emphasizing on trichome type.

## MATERIALS AND METHODS

For this study, specimens with fruits were collected during fieldwork conducted between November 2010 and April 2011, in Santa Catarina State, Brazil. These specimens were deposited at the Herbarium "Barbosa Rodrigues" (HBR) and the Botany Department Herbarium, Federal University of Santa Catarina (FLOR), with abbreviations according to Holmgren, Holmgren, and Barnett (1990). The species that were not found in the field were studied starting from ancient specimens deposited in HBR.

Three fruits of each species were collected from the specimens. The specimens were: *Desmodium adscendens* (Sw.) DC (D. M. Freitas 82 [21/03/2010] FLOR/HBR, 165 [07/02/2011] FLOR/HBR, 179 [17/03/2011] HBR); *D. affine* Schldl. (R. Reitz & R. M. Klein 16752 [01/01/1964] HBR; D. M. Freitas 97 [21/03/2010] HBR, 192 [18/03/2011] FLOR/HBR); *D. barbatum* (L.) Benth. (D. M. Freitas 168 [16/03/2011] HBR, 186 [17/03/2011] FLOR/HBR, 203 [23/04/2011] FLOR/HBR); *D. cuneatum* Hook. & Arn. (Rambo 59229 [03/02/1956] HBR; R. Reitz 5296 [26/01/1953] HBR, 6596 [04/02/19630] HBR); *D. subsecundum* Vogel (R. Reitz & R. M. Klein 3978 [13/04/1956] HBR; A. Reis & D. M. Freitas 2602 [13/02/2011] FLOR/HBR, 2616 [23/02/2011] FLOR/HBR); *D. incanum* DC. (D. M. Freitas 166 [07/02/2011] FLOR/HBR, 187 [18/03/2011] HBR, 204 [10/06/2011] FLOR/HBR); *D. leiocarpum* (Spreng.) G. Don. (A. Reis 2617 [08/03/2011] FLOR/HBR, 2620 [09/02/2011] FLOR/HBR; R. Reitz & R. M. Klein 12294 [22/02/1962]

HBR); *D. pachyrhizum* Vogel (L. B. Smith & R. Reitz 9934 [06/01/1957] HBR; L. B. Smith et al. 14458 [14/01/1965] HBR; R. Reitz 5220 [26/01/1953] HBR); *D. polygaloides* Chodat & Hassl. (R. Reitz 4328 [24/01/1952] HBR; R. Reitz & R. M. Klein 11810 [09/01/1962] HBR, 16579 [29/12/1963] HBR); *D. subsericeum* Malme (R. Reitz & R. M. Klein 8774 [24/04/1959] HBR, 12243 [22/02/1962] HBR; D. M. Freitas & A. Reis 107A [27/05/2010] FLOR/HBR); *D. tortuosum* (Sw.) DC. (D. M. Freitas & A. Reis 145 [12/11/2010] FLOR/HBR; D. M. Freitas & A. Zanin 185 [17/03/2011] FLOR/HBR; A. Reis 2628 [13/03/2011] FLOR/HBR); *D. triarticulatum* Malme (D. M. Freitas 93 [21/03/2010] FLOR/HBR, 200 [22/04/2011] FLOR/HBR; A. Reis 2621 [09/03/2011] FLOR/HBR) and *D. uncinatum* (Jacq.) DC. (D. M. Freitas 195 [22/4/2011] FLOR/HBR; A. Reis 2623 [09/3/2011] FLOR/HBR; R. Reitz & R. M. Klein 14828 [13/4/1963] HBR).

Lima et al. (2014) reported the occurrence of 16 species of *Desmodium* in Santa Catarina State. However, four species noted by these researchers, including *D. album*, *D. arechavaletae*, *D. triflorum* and *D. venosum*, were not found in either HBR or FLOR, nor were they found in the many field collections covering almost every Brazilian state. Meanwhile, in this investigation, the occurrence of *D. leiocarpum* was registered, but not previously reported to Santa Catarina State by Lima et al. (2014).

Images of fruits were captured with a stereoscopic microscope (Olympus SZ-40, Tokyo, Japan) with a digital camera (Sony Cyber-shot™ compact digital camera, 7.2 megapixels, Tokyo, Japan).

Three fruits of each species (total of 39 samples) were rehydrated in distilled water with some drops of detergent and heated to 90°C (Bersier & Bocquet, 1960). SEM analysis was carried out in the Central Laboratory for Electron Microscopy at the Federal University of Santa Catarina – LCME-UFSC. After rehydration, the samples were fixed in glutaraldehyde 2.5%, 0.1M sodium phosphate

buffer, pH 7.2, and then dehydrated in ethanol. Samples were critical point dried with liquid CO<sub>2</sub> (Leica EM model CDP 300, Wetzlar, Germany). Dried samples were placed on aluminum supports with the aid of double-sided carbon tape and coated with 20nm gold in a metallizer (Leica EM model SCD 500, Wetzlar, Germany). The analysis was then carried out at the LCME using SEM (JEOL JSM-6390 LV, Tokyo, Japan).


Thirteen rehydrated samples were examined using light microscopy (LM). Furthermore, two fruits of each species were freehand sectioned transversely and longitudinally, with a razor blade. Sections were exposed to the following histochemical reagents: Sudan III for oils and Thionine for mucilage (Costa, 1982), with the purpose of test presence of chemicals (oil or mucilage) on the trichomes.

## RESULTS

*Desmodium* species have lomentaceous fruits (Fig. 1). While uncinata trichomes were observed, other types can occur, including uniseriate, globose and subulate. Loments of some *Desmodium* species were sinuous at both edges, or the upper edge was straight, while the lower was wavy (Fig. 1). Two species, *D. subsecundum* and *D. leiocarpum*, had glabrescent loments, while the loments of species were pubescent with differential features (Fig. 2). The shape of the articles, presence of distinct types of trichomes and aspects of epidermal cells, with epicuticular ridges and papillae, also proved to be useful in distinguishing among species (Fig. 1, Fig. 2, Fig. 3, Fig. 4, Fig. 5 y Fig. 6).

The structural aspects of trichomes can be classified into uni- or multicellular and may still be simple, i.e., nonglandular or glandular. Five morphological types of trichomes were recognized and could be classified as follows:


1. Uncinate trichome (Fig. 3): glandular, with oil secretion at the base in all species. This trichome type showed dense distribution in *D. adscendens*, *D. affine*, *D. incanum*,




**Fig. 1.** Photomicrographs of loment and articles (bar in loment=5mm/bar in article=2mm). A- *Desmodium adscendens* (Sw.) DC. (Freitas, D. M. 165). B- *D. affine* Schltdl. (Freitas, D. M. 192). C- *D. barbatum* (L.) Benth. (Freitas, D. M. 203). D- *D. subsericeum* Malme (Freitas, D. M. & Reis, A. 107a). E- *D. cuneatum* Hook & Arn. (Reitz, R. 5296). F- *D. incanum* DC. (Freitas, D. M. 187). G- *D. subsecundum* Vogel (Reitz, R. & Klein, R. M. 3978). H- *D. leiocarpum* (Spreng.) G. Don. (Reitz, R. & Klein, R. M. 12294). I- *D. pachyrhizum* Vogel (Reitz, R. 5220). J- *D. polygaloides* Chodat & Hassl. (Reitz, R. & Klein, R. M. 11810). K- *D. tortuosum* (Sw.) DC. (Reis, A. 2628). L- *D. triarticulatum* Malme (Freitas, D. M. 200). M- *D. uncinatum* (Jacq.) DC. (Freitas, D. M. 195).

- D. uncinatum*, *D. subsericeum* and *D. triarticulatum* (Fig. 2A, Fig. 2B, Fig. 2E, Fig. 2I, Fig. 2K and Fig. 2L); however, they were sparse in other species (Fig. 2C, Fig. 2D, Fig. 2F, Fig. 2G, Fig. 2H, Fig. 2J and Fig. 2M).
2. Globular multicellular trichome (Fig. 4A, Fig. 4B, Fig. 4C, Fig. 4D, Fig. 4E, Fig. 4F, Fig. 4G, Fig. 4H, Fig. 4I, Fig. 4J, Fig. 4K): glandular, secretion of oil or mucilage throughout the structure. These trichomes were not found in *D. pachyrhizum* or *D. polygaloides*; however, they were present in all other species and showed a positive reaction to Thionine, indicating the presence of mucilage, except *D. subsecundum* which reacted only with Sudan III for oils.
  3. Globular unicellular trichome (Fig. 4L): glandular, secretion unspecified (non-oily and non-mucilaginous). This trichome type was only present in *D. uncinatum*, and it showed no positive reaction to oil or mucilage. The substance could not be identified in this work.
  4. Uniseriate trichome (Fig. 5): glandular, secretion throughout the structure. This trichome type was absent in *D. subsecundum*. When these trichomes were present, secreted oil was present in *D. pachyrhizum*,





**Fig. 2.** Scanning electromicrographs of the fruit surface (bar=100µm). A- *Desmodium adscendens* (Sw.) DC. (Freitas, D. M. 165). B- *D. affine* Schldtl. (Freitas, D. M. 192). C- *D. barbatum* (L.) Benth. (Freitas, D. M. 203). D- *D. cuneatum* Hook & Arn. (Rambo, B. 59229). E- *D. incanum* DC. (Freitas, D. M. 166). F- *D. leiocarpum* (Spreng.) G. Don. (Reis, A. 2616). G- *D. pachyrhizum* Vogel (Smith, L. B. & Reitz, R. 9934). H- *D. polygaloides* Chodat & Hassl. (Reitz, R. & Klein, R. M. 11810). I- *D. subsericeum* Malme (Freitas, D. M. & Reis, A. 107a). J- *D. tortuosum* (Sw.) DC. (Reis, A. 2628). K- *D. triarticulatum* Malme (Freitas, D. M. 93). L- *D. uncinatum* (Jacq.) DC. (Reitz, R. & Klein, R. M. 14828). M- *D. subsecundum* Vogel (Reis, A. & Freitas, D. M. 2616).




**Fig. 3.** Scanning electromicrographs of uncinete trichomes (bar=50µm). A- *Desmodium adscendens* (Sw.) DC. (Freitas, D. M. 165), arrow: papillary cells. B- *D. affine* Schltdl. (Freitas, D. M. 192), arrow: papillary cells. C- *D. barbatum* (L.) Benth. (Freitas, D. M. 168). D- *D. cuneatum* Hook. & Arn. (Rambo, B. 59229). E- *D. subsecundum* Vogel (Reis, A & Freitas, D. M. 2602), arrow: papillary cells. F- *D. incanum* DC. (Freitas, D. M. 166). G- *D. leiocarpum* (Spreng.) G. Don. (Reis, A. 2620), arrow: epicuticular striations. H- *D. uncinatum* (Jacq.) DC. (Reitz, R. & Klein, R. M. 10802), arrow: epicuticular wax. I- *D. triarticulatum* Malme. (Freitas, D. M. 93), arrow: papillary cells. J- *D. pachyrhizum* Vogel (Smith, L. B. et al. 14458), arrow: epicuticular wax. K- *D. polygaloides* Chodat & Hassl. (Reitz, R. & Klein, R. M. 16579), arrow: epicuticular wax. L- *D. tortuosum* (Sw.) DC. (Freitas, D. M. & Reis, A. 145). M- *D. subsericeum* Malme. (Freitas, D. M. & Reis, A. 107a), arrow: papillary cells.

*D. cuneatum* and *D. polygaloides*, but mucilage was present in *D. adscendens*, *D. affine*, *D. barbatum*, *D. leiocarpum*, *D. incanum*, *D. triarticulatum*, *D. tortuosum*, *D. subsecundum* and *D. uncinatum*.

- Subulate trichome (Fig. 6): glandular, secretion (oil) at the base, when present. This trichome type occurs in *D. barbatum*, *D. pachyrhizum*, *D. cuneatum*, *D. subsecundum*, *D. leiocarpum* and *D. polygaloides*.

Epicuticular wax depositions were found on *D. pachyrhizum* (Fig. 3J, Fig. 5I), *D. polygaloides* (Fig. 3K, Fig. 5G, Fig. 6D), *D. uncinatum* (Fig. 3H) and *D. cuneatum* (Fig. 5H), while *D. leiocarpum* (Fig. 3G, Fig. 5D) showed the presence of epicuticular streaks on the epidermal surface. The epidermal cells in 9 of the 13 species were papillose, characterized by the convexity of the outer periclinal wall, which was more pronounced in *D. affine* (Fig. 3B) and *D. triarticulatum* (Fig. 3I, Fig. 5F), but



**Fig. 4.** Scanning electromicrographs of (A-K) globose multicellular trichomes and (L) globose unicellular trichomes (bar=10µm). A- *Desmodium adscendens* (Sw.) DC. (Freitas, D. M. 179). B- *D. affine* Schltdl. (Freitas, D. M. 97). C- *D. barbatum* (L.) Benth. (Freitas, D. M. 203). D- *D. cuneatum* Hook. & Arn. (Rambo, B. 59229). E- *D. subsecundum* Vogel (Reis, A. & Freitas, D. M. 2616). F- *D. incanum* DC. (Freitas, D. M. 204). G- *D. leiocarpum* (Spreng.) G. Don. (Reis, A. 2617). H- *D. subsericeum* Malme (Freitas, D. M. & Reis, A. 107a). I- *D. triarticulatum* Malme (Reis, A. 2621). J-L- *D. uncinatum* (Jacq.) DC. (Reitz, R. & Klein, R. M. 14828), J- globose multicellular trichomes, L- arrow: globose unicellular trichome. K- *D. tortuosum* (Sw.) DC. (Reis, A. 2628).

also present in *D. adscendens* (Fig. 3A), *D. barbatum* (Fig. 5C), *D. subsecundum* (Fig. 3E), *D. leiocarpum* (Fig. 5D, Fig. 6E), *D. subsericeum*


(Fig. 3M, Fig. 5K) and *D. tortuosum* (Fig. 5J, Fig. 6C). In fruits of *Desmodium*, stomata were found on the fruit surface (Fig. 5B).

**Key for identification of *Desmodium* species in Santa Catarina, Brazil, using fruit macro- and micromorphology**

1. Loments with both edges wavy .....	2
1. Loments with top edge straight and bottom edge wavy .....	9
2. Loments glabrescent .....	3
2. Loments densely pilous .....	4
3. Loments with orbicular articles, without uniseriate trichomes .....	<i>D. subsecundum</i>
3. Loments with elliptical or ovate articles, with uniseriate trichomes .....	<i>D. leiocarpum</i>
4. Loments with twisted articles .....	<i>D. tortuosum</i>
4. Loments without twisted articles .....	5
5. Loments with articles triangular or subtriangular, without trichomes subulate .....	6
5. Loments with articles elliptical, orbicular or suborbicular; trichomes subulate type .....	7
6. Articles subtriangulate, ca. 3-5 x 2-3mm, epidermal cells slightly papillose .....	<i>D. subsericeum</i>
6. Articles triangulate, ca. 6-7 x 4mm; epidermal cells strongly papillose .....	<i>D. triarticulatum</i>
7. Loments with isthmuses subcentral, articles suborbicular, 4-6 x 3-5mm, globular multicellular trichome type .....	<i>D. cuneatum</i>
7. Loments with isthmuses central, articles elliptical or orbicular, 3-5 x 1-3mm, without globular multicellular trichomes .....	8




8. Articles orbicular and elliptical, ca. 3 x 1-2mm, subcoriaceous ..... *D. pachyrhizum*  
 8. Articles orbicular, ca. 4 x 2-3mm, membranaceous ..... *D. polygaloides*  
 9. Articles with sparse uncinata trichomes, with subulate trichomes ..... *D. barbatum*  
 9. Articles with abundant uncinata trichomes, without subulate trichomes ..... 10  
 10. Articles triangular, with globose unicellular trichomes and epicuticular wax depositions ..... *D. uncinatum*  
 10. Articles square, ovate-depressed or round-depressed, without globose unicellular trichomes and epicuticular wax depositions ..... 11  
 11. Articles quadrangular, 4-6 x 3-5mm, leathery ..... *D. incanum*  
 11. Articles ovate-depressed or rounded-depressed, 5-7 x 3-4mm ..... 12  
 12. Articles ovate-depressed with epidermal cells strongly papillose ..... *D. affine*  
 12. Articles rounded-depressed; epidermal cells slightly papillose ..... *D. adscendens*



**Fig. 5.** Scanning electromicrographs of uniseriate trichomes (bar=50µm). A- *Desmodium adscendens* (Sw.) DC. (Freitas, D. M. 165). B- *D. affine* Schltld. (Freitas, D. M. 97), arrow: stoma. C- *D. barbatum* (L.) Benth. (Freitas, D. M. 203), arrow: papillary cells. D- *D. leiocarpum* (Spreng.) G. Don. (Reis, A. 2620), arrow: epicuticular striations and papillary cells. E- *D. incanum* DC. (Freitas, D. M. 166). F- *D. triarticulatum* Malme (Freitas, D. M. 93), arrow: papillary cells. G- *D. polygaloides* Chodat & Hassl. (Reitz, R. & Klein, R. M. 16579), arrow: epicuticular wax. H- *D. cuneatum* Hook. & Arn. (Rambo, B. 59229), arrow: epicuticular striations. I- *D. pachyrhizum* Vogel (Smith, L. B. et al. 14458), arrow: epicuticular wax. J- *D. tortuosum* (Sw.) DC. (Freitas, D. M. & Reis, A. 145), arrow: papillary cells. K- *D. subsericeum* Malme (Freitas, D. M. & Reis, A. 107a), arrow: papillary cells. L- *D. uncinatum* (Jacq.) DC. (Reitz, R. & Klein, R. M. 14828), arrow: globose unicellular trichome.





**Fig. 6.** Scanning electromicrographs of subulate trichomes (bar=50 $\mu$ m). A- *D. barbatum* (L.) Benth. (Freitas, D. M. 186). B- *D. cuneatum* Hook. & Arn. (Reitz, R. 5296). C- *D. tortuosum* (Sw.) DC. (Freitas, D. M. & Reis, A. 185), arrow: papillary cells. D- *D. polygaloides* Chodat & Hassl. (Reitz, R. & Klein, R. M. 16579), arrow: epicuticular wax. E- *D. leiocarpum* (Spreng.) G. Don. (Reis, A. 2617), arrow: papillary cells. F- *D. subsecundum* Vogel (Reitz, R. & Klein, R. M. 3978). G- *D. pachyrhizum* Vogel (Smith, L. B. et al. 14458).

## DISCUSSION

The presence of trichomes on plants, as noted by Werker (2000), may serve as a mechanical barrier against various external factors, such as herbivores, pathogens, intense light, and temperature extremes. In the case of glandular trichomes, these structures can provide chemical protection to the plant by secreting substances, such as mucilage, resins or lipids, which may poison or deter herbivores and pathogens. Trichomes may also aid in dispersion, using, for example, bristles or hooks attached to the fur and feathers of animals.

Uncinate trichomes were present in all species analyzed. Such hook-shaped trichomes play an important role in the dispersion of the diaspore by promoting fixation to the skin of animals and the clothing of humans. However, by comparing these species, it was observed that the length of trichomes is distinctly variable in different species, and that the distribution of these structures on the surface of the fruit can be dense or sparse, also depending on the species. While significant differences can be observed among some trichomes, homogeneity is seen among others. From these results, it can be concluded that dispersal of fruit in *D. adscendens*, *D. affine*, *D. incanum*, *D. subsericeum*, *D. triarticulatum* and *D. uncinatum* is epizoochoric since uncinata trichomes were more densely distributed on the fruit surface. However, in *D. barbatum*, *D. cuneatum*, *D. pachyrhizum*, *D. polygaloides* and *D. tortuosum*, the uncinata trichomes are sparse, indicating two forms of fruit dispersal: epizoochoric and anemochoric. For *D. subsecundum* and *D. leiocarpum*, uncinata trichomes are short and sparsely distributed on the fruit surface, suggesting the anemochoric type. As already mentioned by Barroso, Morim, Peixoto, and Ichaso (1999), most species of *Desmodium* present articles covered with uncinata trichomes, which make them suitable for epizoochoric dispersion; however, anemochoric dispersion is also observed in the genus. Moreover, according to Lima (2011), the presence of uncinata trichomes is uncommon in fruits of all species,

and possibly, the loss of such trichomes is associated with changes in dispersal strategies.

Globular multicellular trichomes were small in size and their presence could only be detected by optical or scanning electronic microscopy. These trichomes were present in 11 of the 13 species examined. Globose unicellular trichomes were only found on *D. uncinatum*.

Other surface features were observed in the epidermal ultrastructural examination of the fruit surfaces. Specifically, for example, epidermal cells were papillose in nine of the 13 species. While this feature can be used to help identify a given species, it should be done with caution because it may be in which such species occurs. The convexity of the outer walls of the epidermal cells, according Larcher (2000), plays an important role in xeric environments, reflecting solar radiation, while minimizing the effects of high solar irradiation on internal tissues.

Epicuticular wax depositions were found on three species, including *D. pachyrhizum*, *D. polygaloides* and *D. uncinatum*. These depositions also reflect environmental conditions, with the greatest wax deposits occurring with higher exposure to solar radiation (Cutter, 1978; Barthlott, 1990). However, the type of deposition may vary and has been considered a taxonomic character (Barthlott et al., 1998). Furthermore, *D. cuneatum* and *D. leiocarpum* showed epicuticular streaks on the epidermal surface, which is also related to xeric environments and may, therefore, be complementary to the characterization of the species, but not a determinant of identification. According to Monteiro, Castro, and Giulietti (1985), the presence of epicuticular ornamentation can provide protection for the stomata or increase the reflection of sunlight. In fruits of *Desmodium*, the presence of stomata may promote evaporation to maintain homeostasis of internal water supply in environments exposed to excessive solar irradiation.

The data on macro- and micromorphological characteristics of the fruit enabled the development of an identification key of

*Desmodium* species in Santa Catarina, Brazil. The following features were shown to be useful for the design of this key: form of the loment margin, form of article, indument glabrescent or pilous, trichome type, papillous epidermal cells present or not, and epicuticular striations present or not.

#### ACKNOWLEDGMENTS

We thank the administrators the Herbarium "Barbosa Rodrigues" (HBR) by loan of specimens used in this study and support from Deise Rebelo Consoni for SEM analysis in Central Laboratory for Electron Microscopy at the Federal University of Santa Catarina –LCME-UFSC, Brazil.

#### RESUMEN

Características morfológicas y micromorfológicas de frutos *Desmodium* (Leguminosae: Papilionoideae). El género *Desmodium* está representado en Santa Catarina, Brasil, por 13 especies, todas con frutos lomentaceos. Los lomentos han tenido variación en forma, tamaño y características del margen del istmo, y la superficie es glabra o cubierta por tricomas de diferentes tipos. La diversidad morfológica de los tricomas se vuelve particularmente relevante para la descripción taxonómica. Los tipos de tricomas presentes en la superficie de los frutos *Desmodium*, proporcionan datos para la identificación y clasificación de las especies en el Estado de Santa Catarina, Brasil. Tres frutos de cada especie fueron recogidos y depositados en el Herbario (HBR y FLOR, Santa Catarina, Brasil). Algunas muestras fueron rehidratadas y examinadas usando microscopía de luz (LM), las secciones fueron expuestas a los siguientes reactivos histoquímicos: Sudan III para los aceites y Tionina para mucílago. Los aspectos estructurales de los tricomas se pueden clasificar en uni o multicelulares y pueden todavía ser simples, es decir, no glandular o glandular. Por medio del uso de la microscopía electrónica de barrido (SEM), cinco tipos de tricomas se han identificado y analizado entre las especies de *Desmodium* estudiadas: uncinado, uniseriado, globoso multicelular, globoso unicelular y subulado. Características como el margen del lomento y la forma del artículo, indumento glabrescente o piloso, tipo de tricomas con o sin células epidérmicas papilosas, y estrías epicuticulares mostraron

valor de diagnóstico relevante. La clave de identificación fue desarrollada para especies de *Desmodium* del Estado de Santa Catarina, Brasil, basado en caracteres macro y micromorfológicos del fruto.

**Palabras clave:** dispersión, clave de identificación, microscopía electrónica de barrido (SEM), estructuras secretoras, taxonomía, tricomas.

#### REFERENCES

- Agbagwa, I. O. & Okoli, B. E. (2005). Fruit Epidermal Micromorphology in the Systematics of *Abrus* Adanson (Papilionaceae) in Parts of Tropical West Africa. *Asian Journal of Plant Sciences*, 4(6), 652-659.
- Akcin, O. E. (2008). Seed coat and fruit surface micromorphology of some *Cynoglossum* L. (Boraginaceae) species. *Bangladesh Journal of Botany*, 37(2), 115-119.
- Ávalos, S. V. & Salinas, A. D. (2003). Los tricomas foliares en la caracterización de un grupo de especies del género *Quercus*, sección *Lobatae* (Fagaceae). *Anales del Instituto de Biología, Universidad Nacional Autónoma de México, Serie Botánica*, 74(1), 5-15.
- Azevedo, M. G. (1981). *O gênero Desmodium Desv. no Brasil: considerações taxonômicas* (Master's thesis, Universidade Estadual de Campinas, Brasil). Retrieved from <http://www.bibliotecadigital.unicamp.br/document/?code=vtls000075165>
- Barroso, G. M., Morim, M. P., Peixoto, A. L., & Ichaso, C. L. F. (1999). *Frutos e sementes - morfologia aplicada à sistemática de dicotiledóneas*. Viçosa: Editora UFV.
- Barthlott, W. (1990). Scanning electron microscopy of the epidermal surface in plants. In D. Claugher (Ed.), *Scanning Electron Microscopy in Taxonomy and Functional Morphology* (pp. 69-94). Oxford: Clarendon Press.
- Barthlott, W., Neinhuis, C., Cutler, D., Ditsch, F., Meusel, L., Theisen, I., & Wilhelmi, H. (1998). Classification and terminology of plant epicuticular waxes. *Botanical Journal of the Linnean Society*, 126(3), 237-260.
- Bersier, J. D. & Bocquet, G. (1960). Les méthodes d'éclaircissement en vascularisation et en morphogénie végétales comparés. *Archives des Sciences*, 13, 555-566.
- Costa, A. F. (1982). *Farmacognosia: farmacognosia experimental*. Lisboa: Fundação Calouste Gulbenkian.
- Cutter, E. G. (1978). *Plant anatomy*. London: Edward Arnold Publishers.



- Fahn, A. & Cutler, D. F. (1992). *Xerophytes*. Germany: Gebrüder Borntraeger.
- Holmgren, P. K., Holmgren, N. H., & Barnett, L. C. (1990). *Index Herbariorum Part I: The Herbaria of the World*. New York: New York Botanical Garden.
- Kaya, A., Ünal, M., Özgökçe, F., Doğan, B., & Martin, E. (2011). Fruit and seed morphology of six species previously placed in *Malcolmia* (Brassicaceae) in Turkey and their taxonomic value. *Turkish Journal of Botany*, 35, 653-662.
- Larcher, W. (2000). *Ecofisiologia Vegetal*. São Carlos: Editora Rima.
- Lima, L. C. P., Oliveira, M. L. A. A., & Tozzi, A. M. G. A. (2014). *Desmodium* Desv. In *Lista de Espécies da Flora do Brasil*. Rio de Janeiro: Jardim Botânico do Rio de Janeiro. Retrieved from <http://floradobrasil.jbrj.gov.br/jabot/floradobrasil/FB22930>
- Lima, L. C. P. (2011). *Estudos Filogenéticos em Desmodium Desv. (Leguminosae-Papilionoideae) na América do Sul e Revisão Taxonômica das Espécies Brasileiras* (Doctoral Dissertation, Universidade Estadual de Feira de Santana, Brasil). Retrieved from [http://www2.uefs.br/ppgbot/pdf\\_dissertacoes\\_teses/doutorado/2011/TeseFinal\\_LauraLima.pdf](http://www2.uefs.br/ppgbot/pdf_dissertacoes_teses/doutorado/2011/TeseFinal_LauraLima.pdf)
- Monteiro, W. R., Castro, M. M., & Giulietti, A. M. (1985). Aspects of leaf structure of some species of *Leiotrix* Ruhl. (Eriocaulaceae) from Serra do Cipó (Minas Gerais, Brasil). *Revista Brasileira de Botânica*, 7, 137-147.
- Ohashi, H. (2014). Tribe Desmodieae. In G. P. Lewis, B. Schrire, B. Mackinder, & M. Lock (Eds.), *Legumes of the World*. Kew: Royal Botanic Gardens.
- Oliveira, M. L. A. A. (1983). Estudo Taxonômico do gênero *Desmodium* Desv. (Leguminosae, Faboideae, Desmodieae). *Iheringia, Série Botânica*, 31, 37-104.
- Oliveira, M. L. A. A. (1990). Adições para o gênero *Desmodium* Desvaux (Leguminosae-Faboideae) no Rio Grande do Sul, Brasil. *Iheringia, Série Botânica*, 40, 77-87.
- Ritter, M. R. & Miotto, S. T. S. (2006). Micromorfologia da superfície do fruto de espécies de *Mikania* Willd. (Asteraceae) ocorrentes no Estado do Rio Grande do Sul, Brasil. *Acta Botanica Brasílica*, 20, 241-247.
- Setubal, R. B., Lima, L. C. P., & Grings, M. (2010). Espécie campestre provavelmente extinta (*Desmodium craspediferum* Azevedo & Oliveira, Fabaceae) reencontrada no Rio Grande do Sul, Brasil. *Revista Brasileira de Biociência*, 8, 432-438.
- Werker, E. (2000). Trichome Diversity and Development. In D. L. Hallahan & J. C. Gray (Eds.), *Advances in Botanical Research Incorporating Advances in Plant Pathology: Plant Trichomes* (Vol. 31, pp. 1-35). London: Academic Press.