

Water pollution and distribution of the black fly (Diptera: Simuliidae) in the Atlantic Forest, Brazil

Tatiana N. Docile^{1,2,3}, Ronaldo Figueiró^{3,4}, Leonardo H. Gil-Azevedo⁵ & Jorge L. Nessimian¹

1. Laboratório de Entomologia, Departamento de Zoologia, Universidade Federal do Rio de Janeiro (UFRJ), Av. Carlos Chagas Filho, 373, Edifício do Centro de Ciências da Saúde, Bloco A, sala A1-107, Ilha do Fundão, Rio de Janeiro, RJ, Brasil – CEP: 21941-902; Caixa Postal: 68044; tatidocile@gmail.com; jnessimian@gmail.com
2. Programa de Pós-Graduação em Ecologia da Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil.
3. Laboratório de Biotecnologia Ambiental, Centro Universitário Estadual da Zona Oeste (UEZO), Rio de Janeiro, Brazil; ronaldofigueiro@gmail.com
4. Centro Universitário de Volta Redonda (UniFOA), Volta Redonda, Brazil.
5. Departamento de Entomologia, Museu Nacional, Universidade Federal do Rio de Janeiro (UFRJ) Rio de Janeiro, Brazil; lhgazevedo@gmail.com

Received 03-X-2014. Corrected 25-II-2015. Accepted 27-III-2015.

Abstract: Black flies have medical importance because some species are vectors of the Onchocerciasis and Mansonellosis, nevertheless, their ecology and potential use as bioindicators is still poorly studied in the Neotropical Region. In Brazil, bioindicators use is strongly focused in a multimetric ecological index approach; this way, we investigated the black fly spatial distribution, in relation to abiotic factors correlated to water quality, to provide baseline information for their utilization as standalone indicators of lotic systems integrity. We have tested the hypothesis that environmental changes related to urbanization, lead to decreased abundance and loss in the number of species of the black fly fauna. The sampling was conducted in 10 urban and 10 preserved streams during the dry season (August-September) of 2012, in the mountainous region of Teresópolis, State of Rio de Janeiro, Brazil. The streams were characterized for their environmental integrity conditions and physico-chemical properties of water. In each stream, five different rapid points were sampled in a section of 50 meters, 10 meters apart from each other. The black flies were sampled with a kick-net sampler on rocky substrates. The material was separated and the larvae were sorted in morphotypes, and later, the final instar specimens were dissected and identified with the help of taxonomical literature at species level. A total abundance of 488 larvae from nine species were collected, 5 (1.02 %) in extremely impacted streams, 470 (96.31 %) in intermediate streams and 13 (2.66 %) in preserved streams. The visual evaluation (HII) differed in relation to the water physico-chemical evaluation, in which more variation in the characterization of the sampling sites was observed. In Canonical Correspondence Analysis *Simulium subpallidum*, *S. inscrustatum* and *S. pertinax* were significantly associated with intermediate values of most of the variables, and then to intermediate impacted sites. On the other hand, *Lutzsimulium hirticosta*, *S. subnigrum* and *Simulium* sp. A were associated to low values of chemical variables, and then to more preserved sites. Most studies on the bioindicator potential of Simuliidae have suggested an approach to agricultural impacts, while our results, on the other hand, showed that Simuliidae species were present in streams with intermediate urban pollution impacts, but absent in heavily impacted sites. Thus, our data suggested that some species are associated to more pristine breeding sites, such as *L. hirticosta* and *Simulium* sp. A, while others may be good bioindicators of moderately impacted streams, such as *S. pertinax*, *S. subnigrum* and *S. subpallidum*. Rev. Biol. Trop. 63 (3): 683-693. Epub 2015 September 01.

Key words: black flies, community ecology, environmental characterization, impacted streams, tropical forest.

Aquatic ecosystems, such as streams, lakes, rivers and reservoirs have been disturbed and significantly affected by increasing anthropic activities (McAllister, Hamilton, &

Harvey, 1997). The most serious problems are seen in urban areas where watercourses receive input of industrial and domestic sewages, sediment and trash, changes such as plumbing and



waterproofing of soils, resulting in the modification of its physical, chemical and biological structure (Shepp & Cummins, 1997). There is a change of abiotic environment due to various impacts suffered, and consequently, there is a decrease in water quality, loss of aquatic biodiversity and changes in the dynamics of biological communities (Goulart & Callisto, 2003; Gutiérrez-Fonseca & Lorion, 2014; Shepp & Cummins, 1997).

Different organisms are used in evaluations of environmental impacts on aquatic ecosystems, including macroinvertebrates (especially insects): these may be viable for laboratory studies when their ecological characteristics are known, and can respond more quickly to changes in the environment, through changes in the community structure (Callisto, Moretti, & Goulart, 2001; Goulart & Callisto, 2003; Reece & Richardson, 1999). The pattern of distribution of aquatic organisms is the result of these biotic and abiotic interactions, the habitat (substrate, flow, turbulence, pH, among others) and food availability (Resh & Rosenberg, 1984). Communities respond to different environmental factors in different ways, including changes in the composition, structure and functional characteristics of the species (Fenoglio, Badino, & Bona, 2002; Heino, Muotka, & Paavola, 2003).

Black flies are among the most abundant components of benthos in lotic systems. This family comprises 2 151 valid species (Adler & Crosskey, 2014). Simuliidae has a worldwide distribution, which is locally limited by the existence of proper lotic microhabitats for their immatures (Eymann, 1993), trophic relationships (McIntosh & Townsend, 1996), as well as competition (Colbo & Porter, 1979) and food availability (Hart, 1986).

Simuliidae is an important family as some of its species are vectors of the Onchocerciasis and Mansonellosis (Cerqueira, 1959; Shelley, Hernández, Maia-Herzog, Dias, & Garritano, 2010). In addition, bites of females cause allergies to humans and certain animals, causing losses in agriculture and tourism (Couceiro, Hamada, Sagot, & Pepinelli, 2014). Thus, this

family has an economic and medical importance (Shelley et al., 2010).

Simuliidae ecology and their potential as bioindicators are still poorly studied in the Neotropical Region (Figueiró & Gil-Azevedo, 2010; Travi & Vargas, 1978). Particularly in Brazil, most of the few recent studies on behavior of black fly immatures is restricted to the Amazon (e.g. Hamada, McCreadie, & Adler, 2002), Mata Atlântica biomes (e.g.: Bertazo & Figueiró, 2012; Coppo & Lopes, 2010; Couceiro et al., 2014; Figueiró, Araújo-Coutinho, Gil-Azevedo, Nascimento, & Monteiro, 2006; Figueiró, Nascimento, Gil-Azevedo, Maia-Herzog, & Monteiro, 2008; Pepinelli, Trivinho-Strixino, & Hamada, 2005;), and just a few in the Cerrado (Figueiró et al. 2015; Figueiró, Gil-Azevedo, Maia-Herzog, & Monteiro, 2012; Figueiró, Maia-Herzog, Gil-Azevedo, Monteiro, 2014; Landeiro, Pepinelli, & Hamada, 2009).

Even though the bioindicator potential of black flies is a concept well established in the literature, only a few recent studies deal with the actual effect of habitat quality in the distribution of black fly larvae in the Neotropical region (e.g.: Strieder, Ronchi, Stenert, Scherer, & Neiss, 2006) and Palearctic regions (e.g.: Illesova, Beracko, Krno, & Halgos, 2010; Kazanci & Ertunc, 2010).

The use of bioindicators in Brazil is strongly focused in a multimetric ecological index approach (Docile & Figueiró, 2013). In the present study we have investigated the black fly spatial distribution in relation to abiotic factors correlated to water quality, in order to provide data that may allow its use as a standalone indicator of lotic systems integrity. In this sense, we have tested the hypothesis that environmental changes, related to urbanization, lead to decreased abundance and loss of the black fly fauna number of species.

MATERIALS AND METHODS

Study area and sampling sites: The study was conducted in the River Watershed Paquequer located in the municipality of Teresópolis,

mountainous region of the State of Rio de Janeiro, (coordinates 22°28'35" S - 43°03'19" W - 22°12'56" S - 42°55'00" W), and covering an area of 12 904 km². In this region, the rainy season occurs from November to February (more than 250 mm/month), and the dry season from June to September (less than 100 mm/month). The natural vegetation in the region is part of the Atlantic Forest biome. Population profile is predominantly urban (96.3 %) and

is concentrated in the district headquarters, Teresópolis. Twenty streams of first and second order were chosen for sampling: 10 outside the urban perimeter and other 10 inside the urban perimeter of Teresópolis (Table 1). The sampling was conducted during the dry season, August and September 2012.

Abiotic factors: The sampling sites were classified in relation to their integrity degree

TABLE 1
Location, altitude, width, order, value of Habitat Integrity Index and characterization of selected for collection in Teresópolis, RJ streams

Sites	Classification	Latitud N	Longitud W	Alt. (m)	Width (m)	Order	HII	Characterization
1	Urban	22°26'26.31"	42°59'8.03"	943	4.0	2 ^a	0.5	On private property; houses nearby; little riparian vegetation; household waste; little silted
2	Urban	22°26'13.76"	42°58'56.17"	898	3.0	1 ^a	0.5	Located in the city street; nearby dwellings; much trash; riparian vegetation; quite silted
3	Urban	22°26'02.36"	42°59'03.50"	954	8.0	2 ^a	0.5	Houses Nearby, presence of trash, with riparian vegetation, some substrates
4	Urban	22°27'22.16"	42°58'53.79"	899	6.0	1 ^a	0.4	In residential condominium; little waste, little riparian vegetation; cemented stretch
5	Urban	22°27'25.60"	42°57'53.80"	900	9.5	1 ^a	0.5	In residential condominium; trash; riparian vegetation; portions output of domestic sewage
6	Urban	22°26'13.12"	42°57'57.95"	952	2.0	1 ^a	0.4	Located across the street; houses nearby, piped stretch; much household waste
7	Urban	22°26'07.56"	42°56'47.94"	920	5.0	1 ^a	0.4	Located across the street; houses nearby; much household waste; excavations and siltation; without retention walls
8	Urban	22°26'05.88"	42°56'28.50"	916	2.5	1 ^a	0.4	On private property; nearby dwellings; little riparian vegetation; much rubbish; cemented excerpts
9	Urban	22°24'50.44"	42°57'01.00"	882	2.0	2 ^a	0.4	Located across the street from the gas station, piped and cemented; few substrates; inlet chemicals category, lack of housing upstream
10	Urban	22°25'20.55"	42°59'03.54"	869	1.0	2 ^a	0.4	Located across the street, piped and cemented; few houses nearby substrates; much trash
11	Preserved	22°26'28.94"	42°59'44.49"	1 125	2.8	1 ^a	0.9	On private property near the PARNASO; neighborhood away from the city; much riparian vegetation; many substrates
12	Preserved	22°27'11.54"	42°59'40.52"	1 058	9.0	1 ^a	0.9	On private property near the PARNASO; neighborhood away from the city; much riparian vegetation; many substrates
13	Preserved	22°27'13.91"	42°59'35.08"	1 074	4.0	1 ^a	0.9	On private property near the PARNASO; neighborhood away from the city; much riparian vegetation; many substrates

TABLE 1 (Continued)

Sites	Classification	Latitud N	Longitud W	Alt. (m)	Width (m)	Order	HII	Characterization
14	Preserved	22°27'23.65"	42°59'50.61"	1 036	6.0	1 ^a	1.0	In PARNASO; much riparian vegetation; many substrates, lack of housing; preservation area
15	Preserved	22°27'19.05"	42°59'56.92"	1 152	3.0	1 ^a	1.0	In PARNASO; much riparian vegetation; many substrates, lack of housing; preservation area
16	Preserved	22°26'50.03"	43°00'20.53"	1 228	3.5	1 ^a	1.0	In PARNASO; access by trail; much riparian vegetation; many substrates, lack of housing; preservation area
17	Preserved	22°26'53.80"	43°00'27.40"	1 206	5.0	2 ^a	1.0	Stream Beija-Flor, in PARNASO; access by trail; much riparian vegetation; many substrates, lack of housing; preservation area
18	Preserved	22°27'02.11"	42°56'32.15"	1 035	3.0	1 ^a	1.0	In the preservation area ("Núcleo Vale da Revolta"); much riparian vegetation; many substrates, lack of housing
19	Preserved	22°27'02.97"	42°56'31.15"	1 035	2.5	1 ^a	1.0	In the preservation area ("Núcleo Vale da Revolta"); much riparian vegetation; many substrates, lack of housing
20	Preserved	22°26'55.46"	42°56'59.16"	1 016	1.0	1 ^a	1.0	In the preservation area ("Núcleo Vale da Revolta"); near concierge; much riparian vegetation; many substrates, lack of housing

using a visual protocol based on Petersen (1992) and Barbour, Gerritsen, Snyder and Stribling (1999) to produce a habitat integrity index (HII) as performed in Nessimian et al. (2008). This protocol is composed of 31 items describing stream environmental condition in relation to land use, riparian zone, streambed characteristics, and stream channel morphology, aquatic vegetation, presence of domestic or industrial effluents and other urban-related characteristics. Each item is composed of five alternatives ordered in a way to represent a degree of integrity. The HII value varies from zero to one, the closer the value is to 1, the more integrate the system is.

Fifteen physical and chemical water variables were measured. At the time of sampling, water temperature and dissolved oxygen (mg O₂/L) were measured using a portable oxymeter (YSI-model F-1550). A sample of water (2 L) was removed from each stream and transported at 12 °C, to posterior laboratory analyses of the following variables: turbidity (HI 93414); electrical conductivity and salinity (YSI-model EC 300 condutivimeter); pH and

alcalinity (Camourze, 1994); total phosphorous, dissolved phosphorous and orthophosphate through the method proposed by Mackereth, Heron and Talling (1978); ammonia (Bower & Holm-Hansen, 1980); total nitrogen and nitrate through the Kejdahl method described in Mackereth et al. (1978); chlorophyll *a* in the watercolumn and periphyton through the method described in Nusch and Palme (1975). The negative values and zero means that the abiotic variable was not detected in the analysis. These analyses were made in the Laboratório de Limnologia, Instituto de Biologia, UFRJ.

Larvae sampling and identification: The black flies were sampled on rocky substrates in riffle /rapids in a section of 50 m along the stream with the help of a kick-net sampler (area of 30x30 cm) and 500 µm mesh. The samples were placed in plastic bags with 80 % ethanol. The material was separated and the larvae were sorted in morphotypes and later the final instar specimens were identified to species level with the help of taxonomical literature (Coscarón & Coscarón-Arias, 2007; Hamada & Adler,

2001), and by using direct comparison with the material deposited at Museu Nacional Universidade Federal of Rio de Janeiro (MNRJ), where the specimens are currently deposited.

The Pearson correlation coefficient was calculated between all measured variables. For pairs of variables with coefficient above 80 % one of them was removed from subsequent analyzes. The sampling sites were analyzed in relation to their ecological distances using Cluster Analyses (Bray-Curtis distance) using PAST (Hammer, Harper, & Ryan, 2007) statistical package. We removed the sites where there was no presence of Simuliidae to contribute to a higher resolution graphic in the biotic cluster. The relations of the abiotic factors with the spatial distribution of black fly larvae were investigated after log transformation and down-weight of rare species through Canonical Correspondence Analysis using 5000 Monte-Carlo permutations in the CANOCO 4.5 statistical package (Ter Braak & Smilauer, 2002). Subsequently, we performed multiple regressions analysis to investigate if there was a significant relationship of the sampled data and the classification of streams provided by the HII.

RESULTS

The sites 7, 8, 9 and 10 (Table 1) showed the lowest scores of HII (0.4), indicating that

these are the extremely impacted streams, while sites 11 to 20 (Table 1) scored over 0.9, being the most preserved. The variables electrical conductivity, salinity and nitrate showed Pearson's correlation coefficient above 80 % with turbidity (86 %) and total nitrogen (93 %), and were excluded from subsequent analyzes. The visual evaluation (HII) differed in relation to the water physico-chemical evaluation, in which more variation in the characterization of the sampling sites was observed (Table 2). The Cluster Analysis applied to the biotic data (species abundances) grouped sites 8 and 10, impacted streams that showed few black fly individuals. Sites 1 to 5, 9 and 11, considered intermediate, and with great abundance of black flies formed another group, and sites 12 to 15 and 17 (the later three considered preserved), with few individuals formed one group. Black flies were absent in sites which are not grouped in the analysis (Fig. 1a). The abiotic values showed streams 14 to 20 were classified as preserved, while streams 1 to 5, 9 and 11 to 13 were classified as intermediate, and streams 6, 7, 8 and 10 were classified as impacted (Table 2), but the Cluster Analysis grouped 6, 7, 8 and 10, and another group with sites 17, 18, 19 and 20 (Fig. 1b).

A total abundance of 488 larvae were collected, 5 (1.02 %) in extremely impacted streams, 470 (96.31 %) in intermediate streams




Fig. 1. Cluster Analysis (Bray-curtis distance) based on variables of 20 streams in Paquequer river watershed, Teresópolis-RJ (A) biotic; (B) abiotic.

TABLE 2
Values of physicochemical variables measured at each of the points studied in 20 tributaries of the Rio Paqueta, Teresópolis, Rio de Janeiro, Brazil

Sites	T (°C)	DO (mL/L)	pH	Tur (NTU)	TP (µMol/L)	DP (µMol/L)	AM (µMol/L)	ORT (µMol/L)	ALC (mEq/L)	TN (µMol/L)	CLO (µg/L)	CLOS (µg/L)
1	16.7	8.89	5.41	0.48	0.89	1.28	0.03	0.15	111.5	79.75	3.10	199.66
2	16.6	7.17	6.48	0.51	3.08	3.12	-0.04	2.53	163.4	82.99	3.04	561.50
3	17.7	8.66	6.58	0.38	0.51	0.79	-1.05	0.71	198.1	37.08	2.44	2985.70
4	18.1	8.91	6.52	0.38	1.98	2.06	1.31	2.11	305.6	77.82	2.67	799.77
5	16.5	8.88	6.42	0.79	1.18	2.38	9.16	1.01	127.9	36.41	2.28	175.08
6	18	4.75	6.72	2.82	2.65	1.07	63.61	2.19	671.9	27.27	-0.09	20.32
7	17.7	5.43	6.8	10.5	3.01	1.81	10.73	1.72	102.7	510.19	3.31	379.57
8	18	7.5	6.86	12.3	2.15	1.24	46.98	1.08	607.2	277.72	0.44	37.34
9	15.9	8.85	6.76	2.84	1.13	0.3	10.88	0.36	310.5	21.04	2.05	106.28
10	18.7	4.78	6.8	13.1	4.77	3.07	4.94	5.67	194.6	295.55	5.20	0.95
11	17.4	7.93	6.05	0.3	0.17	-278	-2.48	0.13	83.46	12.75	0.46	88.09
12	17.8	8.68	5.74	0.5	2.02	-153	4.67	0.14	183.7	7.35	-0.05	53.99
13	17.8	8.55	6.42	1.21	0.24	-30	16.58	0.14	126.1	39.54	1.17	145.54
14	17.1	9.3	6.3	0.27	0.31	0.23	28.63	0.31	50.1	32.08	0.55	19.40
15	14.2	9.21	6.04	0.73	0.39	0.15	-1.25	0.76	125.8	29.58	0.33	71.20
16	15.6	9.65	6.24	0.5	0.06	0.11	3.78	0.18	58.79	34.42	0.63	15.06
17	16.1	9.55	6.22	0.27	-257	0.58	10.39	0.19	45.3	27.65	-0.38	29.09
18	16.2	8.55	5.29	0.41	-196	0.62	-1.68	0.15	40.02	30.44	1.87	117.32
19	15.1	8.74	6.12	0.26	-135	0.78	-1.58	0.28	44.35	37.33	-0.84	16.65
20	15.7	7.57	5.73	0.39	-174	0.36	0	0.2384	85.73	42.08	0.28	56.12

Rj; T = Temperature, DO = Dissolved Oxygen, Tur = Turbidity, TP = Total Phosphorus, DP = Dissolved Phosphorus; AM = Ammonia, ORT = Orthophosphate; ALC = Alkalinity, TN = Total Nitrogen, CLO = Chlorophyll (water column); CLOS = Chlorophyll (substrate).

and 13 (2.66 %) in preserved streams, distributed in eight species: *Lutzsimulium hirticosta* Lutz, 1909, *S. (Chirostilbia) subpallidum* Lutz, 1910, *S. (Chirostilbia) pertinax* Kollar, 1832, *S. (Inaequalium) subnigrum* Lutz s.l., 1910, *S. (Psaroniocompsa) incrustatum* s.l. Lutz 1910, *Simulium* sp. A, *Simulium* sp. B, *Simulium (Trichodagmia)* and Enderlein 1934. In the extremely impacted streams, only three individuals of *S. pertinax* and two *S. incrustatum* s.l. were found. However, intermediate streams presented all previously listed species, except for *Simulium* sp. B. The most abundant was *S. pertinax* with 235 individuals, followed by *S. incrustatum*, with 113 individuals. On the other hand, abundance and species richness were smaller in preserved streams, where *S. subnigrum* s.l., *S. incrustatum* s.l., *Simulium* sp. A, *Simulium* sp. B and *Lutzsimulium hirticosta* were present. Simuliidae were not found in sites 6, 7, 15, 16, 18 and 19.

The Canonical Correspondence Analysis (Fig. 2) showed that *S. subpallidum* and *S. pertinax* were positively associated to turbidity, total phosphorus and pH, suggesting association to moderately impacted sites. On the other hand, *L. hirticosta* and *Simulium* sp. A were negatively associated to phosphorus, dissolved phosphorus, turbidity, total nitrogen, alkalinity, chlorophylla *a*, orthophosphate, thus indicating association to more preserved sites while *S. subnigrum* and *Simulium* sp. B did not show a clear distributional pattern. *S. incrustatum* seemed to be generalistic. The two axes of the CCA explained 0.798 and 0.623 respectively. The linear regressions of the HII scores with species richness, diversity and abundance were not significant, except for the abundance of *S. pertinax* ($p = 0.01$).

DISCUSSION

Several studies have shown that the presence and abundance of black flies are influenced by various environmental variables, which are directly related to anthropogenic pressures, such as loss of riparian vegetation or frequent disturbances (Feld, Kiel, & Lautenschlager,




Fig. 2. Canonical Correspondence Analysis (CCA) of species distribution in relation to abiotic variables. Species: *S. subn* = *Simulium subnigrum*, sp. A = *Simulium* sp. A, sp. B = *Simulium* sp. B, *L. hirt* = *Lutzsimulium hirticosta*, *S. subp* = *Simulium subpallidum*, *S. incr* = *Simulium incrustatum*, *S. pert* = *Simulium pertinax*. Temp = Temperature, DO = Dissolved Oxygen, Tur = Turbidity, CLOS = Chlorophyll *a* (substrate), TP = Total Phosphorus, TN = Total Nitrogen, ORT = Orthophosphate, ALC = Alkalinity, AM = Ammonia, DP = Dissolved Phosphorus, CLO = Chlorophyll *a* (water column), HII = Habitat Integrity Index.

2002; Rühm, 1998; Timm, 1995; Zhang, Malqvist, & Englund, 1998). Furthermore, black flies are known to react to physical and chemical degradations, including acidification and organic pollution (Seitz, 1992). Black fly larvae are able to live in conditions of anoxia for several hours, and are filter-feeding organisms, feeding on organic matter, which favors their adaptation to different environments.

Another explanation for the results would be the intermediate disturbance hypothesis (Connell, 1978). The intermediate sites were submitted to constant sewage discharges, which seem to generate Figueiró et. al (2014) a moderate mortality in the species, which is not high

enough to the point of them fail to recover, but at the same time, is sufficient to limit growth of more competitive species, which could exclude less competitive species. In this study, urban streams with intermediate disturbance had greater abundance of blackflies. If the low species richness at low intensities of disturbance is due to competitive exclusion, as proposed by this hypothesis, there is something that remains to be tested. stated that even though the IDH has recently met criticism in the literature (Fox, 2013), many authors still defend its validity stating that although it is prone to misinterpretations, resulting from the use of different diversity measures (Sheil & Burslem, 2013), it is still robust when used with species richness (Svensson, Lindegarth, Jonsson, & Pavia, 2012), as in the present study.

The CCA showed that *S. subpallidum*, *S. incrustatum* s.l and *S. pertinax* are positively associated to most variables measured, but the cluster analysis showed their association with intermediate impacts, rather than in more severely impacted sites. Paiva (2000), developed studies in the state of Paraná to observe the survival of some species of black flies under laboratory conditions, with the addition of organic matter in water. His results indicate that organic pollution may increase the species richness of Simuliidae. Additionally, the study of Zhang et al. (1998) reported that certain situations (natural or man-made) can cause an increase in the diversity of the group. The authors argued that communities of black flies can be indicative of disturbed sites, such as, floods, droughts, storms. The Simuliidae are known to deal with perturbations in rivers than many of its competitors and predators (Hemphill & Cooper, 1983).

The relationship of the number of individuals of *S. pertinax* points to the prevalence of this species in locations classified as intermediate streams ($p < 0.05$). This association was also found in another study with *S. pertinax* in streams impacted in countryside (Strieder et al., 2006). It is considered an anthropophilic species, on the Northern coast of São Paulo and Southern Brazil *S. pertinax* is the main species

that exerts a role of insect nuisance due to allergic reactions caused by bites influencing well-being of the population as a whole (Strieder & Corseuil, 1992). Therefore, increasing the number of biological control programs to mitigate these impacts (Araújo-Coutinho, 1995; Regis, Silva, Alice & Santos, 2000). However, these organisms are important to the ecosystem because they represent a bond of connection between particulate matter and predators in the food chain (Malmqvist, 1994).

Results obtained by Strieder et al. (2006) in Southern Brazil, showed higher incidence of *S. pertinax* in streams impacted by high concentrations of nitrite and nitrate in agricultural landscape. Couceiro et al. (2014) also studied the distribution of blackflies in locations with disturbance caused by human modification of the landscape (agricultural practices).

Most studies on the bioindicator potential of Simuliidae have tended to approach agricultural impacts, while our results, on the other hand, suggested that these species are present in intermediate urban pollution impacted streams, but absent in heavily impacted sites. Thus, our data suggested that some species such as *L. hirticosta* and sp. A, are associated to more pristine breeding sites, others, may be bioindicators of moderately impacted streams, such as *S. pertinax*, *S. subnigrum* and *S. subpallidum*.

ACKNOWLEDGMENTS

To Larissa Correa and Clayton Portela (Laboratório de Entomologia, IB, UFRJ) for their help in collecting field, Mario Ribeiro and Claudio Marinho (Laboratório de Limnologia, IB, UFRJ) for their help with water analyses. Instituto Brasileiro do Meio Ambiente e dos Recursos Naturais Renováveis (IBAMA), Instituto Chico Mendes de Conservação da Biodiversidade (ICMBio) and INEA for issuing collecting permits. Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and Fundação Carlos Chagas Filho de Amparo à Pesquisa do Estado do Rio de

Janeiro (FAPERJ) provided financial support by grants and fellowships.

RESUMEN

Contaminación del agua y distribución de la mosca negra (Diptera: Simuliidae) en el bosque Atlántico, Brasil. Aunque las moscas negras tienen importancia médica debido a que algunas de sus especies son vectores de la Oncocercosis y Mansonelosis, su ecología y su potencial como bioindicadores es aún poco estudiado en la Región Neotropical. En Brasil el uso de bioindicadores se centra principalmente en un índice ecológico multimétrico, en este estudio investigamos la distribución espacial de las moscas negras y su relación con los factores abióticos, para su utilización como indicadores de la integridad de los sistemas lóticos. En este sentido, hemos probado la hipótesis de que los cambios ambientales relacionados con la urbanización conducen a la disminución de la abundancia y la pérdida de especies de la mosca negra. El muestreo se llevó a cabo en la región montañosa del Estado de Río de Janeiro, Brasil, en 10 zonas urbanas y 10 arroyos conservados durante la estación seca (agosto-septiembre) de 2012. Los arroyos se caracterizaron por sus condiciones de integridad del ambiente y las propiedades físico-químicas de agua. Las moscas negras se muestrearon en sustratos rocosos con un *kick-net sampler*. El material se separó y las larvas se ordenaron por morfotipos y más tarde los especímenes en último estadio se disectaron e identificaron con la ayuda de la literatura taxonómica a nivel de especie. Una abundancia total de 488 larvas de nueve especies fueron recolectadas, 5 (1.02 %) en arroyos muy afectados, 470 (96.31 %) en intermedios y 13 (2.66 %) en conservados. La evaluación visual (HII) difiere de la físico-química del agua, en la que se observó una mayor variación. En el Análisis Correspondencia Canónica, *Simulium subpallidum*, *S. inscrustatum* y *S. pertinax* estaban asociadas significativamente con sitios de impacto intermedio, para la mayoría de las variables. Por otro lado, *Lutzsimulium hirticosta*, *S. subnigrum* y *Simulium* sp. A se asocian con valores bajos de las variables químicas, relacionado a sitios más conservados. La mayoría de los estudios sobre el potencial bioindicador de Simuliidae tienden a enfocarse en los impactos agrícolas, mientras que nuestros resultados, por otro lado, sugieren que las especies de Simuliidae están presentes en arroyos con impactos intermedios de contaminación urbana, pero ausentes en los sitios fuertemente afectados. Por lo tanto, nuestros datos sugieren que algunas especies tales como *L. hirticosta* y *Simulium* sp. A están asociadas a lugares de reproducción más prístinos, mientras que otras pueden ser bioindicadores de arroyos moderadamente afectados, como *S. pertinax*, *S. subnigrum* y *S. subpallidum*.

Palabras clave: moscas negras, ecología comunitaria, caracterización ambiental, ríos impactados, bosque tropical.

REFERENCES

- Adler, P. H., & Crosskey, R. W. (2014). *World blackflies (Diptera: Simuliidae): a comprehensive revision of the taxonomic and geographical inventory (2014)*. Retrieved from: www.clemson.edu/cafls/departments/esps/biomia/pdfs/blackflyinventory.pdf.
- Araújo-Coutinho, C. J. P. C. (1995). Biological control against simuliids in the State of São Paulo, Brazil. *Memórias do Instituto Oswaldo Cruz*, 90, 131-133.
- Barbour, M. T., Gerritsen, J., Snyder, B. D., & Stribling, J. B. (1999). *Rapid Bioassessment Protocols for Use in Streams and Wadeable Rivers: Periphyton, Benthic Macroinvertebrates and Fish*. Washington, D.C.: Environmental Protection Agency, Office of Water.
- Bertazo, K., & Figueiró, R. (2012). Spatial distribution of black fly (Diptera: Simuliidae) immatures in a water current velocity gradient in Aracruz/ES, Brazil. *Revista Universidade Rural*, 32, 91-101.
- Bower, C. E., & Holm-Hansen, T. (1980). A salicylate-hypochlorite method for determining ammonia in seawater. *Canadian Journal of Fisheries and Aquatic Sciences*, 37, 794-798.
- Callisto, M., Moretti, M., & Goulart, M. (2001). Macroinvertebrados bentónicos como ferramenta para avaliar a saúde de riachos. *Revista Brasileira de Recursos Hídricos*, 6, 71-82.
- Camourze, J. P. (1994). *O metabolismo dos ecossistemas aquáticos: fundamentos teóricos, métodos de estudo e análises*. São Paulo: Editora Edgard, Blucher/Fapesp.
- Cerqueira, N. L. (1959). Sobre a transmissão da Mansonella Ozzardi. *Jornal Brasileiro de Medicina*, 5, 885-914.
- Colbo, M. H., & Porter, G. N. (1979). Effects of the food supply on the life history of Simuliidae (Diptera). *Canadian Journal of Zoology*, 57, 301-306.
- Connell, J. H. (1978). Diversity of tropical rainforests and coral reefs. *Science*, 199, 1304-1310.
- Coppo, T. L., & Lopes, J. (2010). Diversidade de Simuliidae (Diptera: Nematocera) de três cursos d' água no parque ecológico da Klabin S.A. – Telêmaco Borba, Estado do Paraná. *Semina: Ciências Biológicas e da Saúde*, 31, 3-14.
- Coscarón S., & Coscarón-Arias, C. L. (2007). Neotropical Simuliidae (Diptera: Insecta). In J. Adis, J. R. Arias, G. Rueda-Delgado, K. M. Wantzen (Eds.), *Aquatic*



- biodiversity in Latin America* (pp. 685). Sofia, Bulgaria: Pensoft Publishers.
- Couceiro, S., Hamada, N., Sagot, L., & Pepinelli, M. (2014). Black-fly assemblage distribution patterns in streams in disturbed areas in southern Brazil. *Acta Tropica*, *140*, 26-33.
- Docile, T. N., & Figueiró, R. (2013). Histórico e perspectivas da utilização de macroinvertebrados no monitoramento biológico de ecossistemas aquáticos no Brasil. *Acta Scientiae et Technicae*, *1*, 31-44.
- Eymann, M. (1993). Some boundary layer characteristics of microhabitats occupied by larval black flies (Diptera: Simuliidae). *Hydrobiologia*, *259*, 57-67.
- Feld, C. K., Kiel, E., & Lautenschlager, M. (2002). The indication of morphological degradation of streams and rivers using Simuliidae. *Limnologia*, *32*, 273-288.
- Fenoglio, S., Badino, G., & Bona, F. (2002). Benthic macroinvertebrate communities as indicators of river environment quality: an experience in Nicaragua. *Revista Biológica Tropical*, *50*, 1125-1132.
- Figueiró, R., Araújo-Coutinho, C. J. P. C., Gil-Azevedo, L. H., Nascimento, E. S., & Monteiro, R. F. (2006). Spatial and temporal distribution of blackflies (Diptera: Simuliidae) in the Itatiaia National Park, Brazil. *Neotropical Entomology*, *35*, 542-550.
- Figueiró, R., Calvet, A., Gil-Azevedo, L. H., Docile, T. N., Monteiro, R. F., Maia-Herzog, M. (2015). Evidence of phenotypic plasticity of larvae of *Simulium subpallidum* Lutz in different streams from the Brazilian Cerrado. *Revista Brasileira de Entomologia*, *59*, 30-33.
- Figueiró, R., & Gil-Azevedo, L. H. (2010). The role of the Neotropical blackflies (Diptera: Simuliidae) as vectors of the onchocerciasis: a short overview of the ecology behind the disease. *Oecologia Australis*, *14*, 745-755.
- Figueiró, R., Gil-Azevedo, L. H., Maia-Herzog, M., & Monteiro, R. F. (2012). Diversity and microdistribution of black fly (Diptera: Simuliidae) assemblages in the tropical savanna streams of the Brazilian cerrado. *Memórias do Instituto Oswaldo Cruz*, *107*, 362-369.
- Figueiró, R., Nascimento, E. S., Gil-Azevedo, L. H., Maia-Herzog, M., & Monteiro, R. F. (2008). Local distribution of blackfly (Diptera: Simuliidae) larvae in two adjacent streams: the role of water current velocity in the diversity of blackfly larvae. *Revista Brasileira de Entomologia*, *52*, 452-454.
- Figueiró, R., Maia-Herzog, M., Gil-Azevedo, L. H., & Monteiro, R. F. (2014). Seasonal variation in black fly (Diptera: Simuliidae) taxocenoses from the Brazilian Savannah (Tocantins, Brasil). *Journal of Vector Ecology*, *39*, 321-327.
- Fox, J. W. (2013). The intermediate disturbance hypothesis should be abandoned. *Trends in Ecology & Evolution*, *28*, 86-92.
- Goulart, M. D., & Callisto, M. (2003). Bioindicadores de qualidade de água como ferramenta em estudos de impacto ambiental. *Revista FAPAM*, *2*, 78-85.
- Gutiérrez-Fonseca, P. E., & Lorion, C. M. (2014). Application of the BMWP-Costa Rica biotic index in aquatic biomonitoring: sensitivity to collection method and sampling intensity. *Revista de Biología Tropical*, *62*, 275-289.
- Hamada N., & Adler, P. H. (2001). Bionomia e chave para imaturos e adultos de *Simulium* (Diptera: Simuliidae) na Amazônia central, Brasil. *Acta Amazonica*, *31*, 109-132.
- Hamada, N., McCreddie, J. W., & Adler, P. H. (2002). Species richness and spatial distribution of black flies (Diptera: Simuliidae) in streams of central Amazonia, Brazil. *Freshwater Biology*, *47*, 31-40.
- Hammer, O., Harper, D. A. T., & Ryan, P. D. (2007). PAST - Palaeontological Statistics. Retrieved from: http://www.uv.es/~pardomv/pe/2001_1/past/pastprog/past.pdf
- Hart, D. D. (1986). The adaptive significance of territoriality in filter-feeding larval black flies (Diptera: Simuliidae). *Oikos*, *46*, 88-92.
- Heino, J., Muotka, T., & Paavola, R. (2003). Determinants of macroinvertebrate diversity in headwater streams: regional and local influences. *Journal of Animal Ecology*, *72*, 425-434.
- Hemphill, N., & Cooper, S. D. (1983). The effect of physical disturbance of two filter-feeder insects in a small stream. *Oecologia*, *58*, 378-382.
- Illesova, D., Beracko, P., Krno, I., & Halgos, J. (2010). Effects of land use on black fly assemblages (Diptera: Simuliidae) in submontane rivers (West Carpathians, Slovakia). *Biologia*, *65*, 892-898.
- Kazanci, N., & Ertunç, O. (2010). Use of Simuliidae (Insecta, Diptera) species as indicators of aquatic habitat quality of Yeşilirmak River Basin (Turkey). *Review of Hydrobiology*, *3*, 27-36.
- Landeiro, V. L., Pepinelli, M., & Hamada, N. (2009). Species richness and distribution of black flies (Diptera: Simuliidae) in the Chapada Diamantina region, Bahia, Brazil. *Neotropical Entomology*, *38*, 332-339.
- Mackereth, J. F. H., Heron, J., & Talling, J. F. (1978). *Water analysis: some revised methods for limnologists*. Ambleside, Cumbria: Freshwater Biological Association.
- Malmqvist, B. (1994). Preimaginal blackflies (Diptera: Simuliidae) and their predators in a central Scandinavian lake outlet stream. *Annales Zoologici Fennici*, *31*, 245-255.

- Mcallister, D. E., Hamilton, A. L., & Harvey, B. (1997). Global freshwater biodiversity: striving for the integrity of freshwater ecosystems. *Sea Wind*, *11*, 1-142.
- McIntosh, A. R., & Townsend, C. R. (1996). Interactions between fish, grazing invertebrates and algae in a New Zealand stream: a trophic cascade mediated by fish induced changes to grazer behaviour. *Oecologia*, *108*, 174-181.
- Nessimian, J. L., Venticinque, E. M., Zuanon, J., De Marco, P. JR., Gordo, M., Fidelis, L., Batista, J. D., & Juen, L. (2008). Land use, habitat integrity, and aquatic insect assemblages in Central Amazonian streams. *Hydrobiologia*, *614*, 117-131.
- Nusch, E. A., & Palme, G. (1975). Biologische methoden für die praxis der gewässeruntersuchung. Bestimmung des Chlorophyll a und phaeopigmentgehaltes in oberflächenwasser. *GWF-Wasser/Abwasser*, *116*, 562-565.
- Paiva, D. P. (2000). *O borrachudo como indicador biológico da qualidade da água*. Concórdia, SC: Embrapa Suínos e Aves.
- Pepinelli, M., Trivinho-Strixino, S., & Hamada, N. (2005). Imaturos de Simuliidae (Diptera: Nematocera) e caracterização de seus criadouros no Parque Estadual Intervales, SP, Brasil. *Revista Brasileira de Entomologia*, *49*, 527-530.
- Petersen, R. C. J. R. (1992). The RCE: a riparian, channel, and environmental inventory for small streams in agricultural landscape. *Freshwater Biology*, *27*, 295-306.
- Reece, P. F., & Richardson, J. S. (1999). Biomonitoring with the reference condition approach for the detection of aquatic ecosystems at risk. In L. M. Darling (Ed.), *Proceedings of a Conference on the Biology and Management of species and habitats at risk* (pp. 549-552). Retrieved from: <http://www.env.gov.bc.ca/wld/documents/re11reece.pdf>
- Regis, L., Silva, S. B., Alice, M., & Santos, M. (2000). The Use of Bacterial Larvicides in Mosquito and Black Fly Control Programmes in Brazil. *Memórias do Instituto Oswaldo Cruz*, *95*, 207-210.
- Resh, V., & Rosenberg, D. (1984). *The ecology of aquatic insects*. New York: Praeger.
- Rühm, W. (1998). Das Habitat und seine Strukturen als Voraussetzung für die autochthone Entwicklung von Schadgebieten der Kriebelmuücken (Diptera: Simuliidae). *Entomologia Generalis*, *23*, 27-37.
- Seitz, G. (1992). Verbreitung und Ökologie der Kriebelmuücken (Diptera: Simuliidae) in Niederbayern. *Lauterbornia*, *11*, 1-231.
- Sheil, D., & Burslem, D. F. R. P. (2013). Defining and defending Connell's intermediate disturbance hypothesis: a response to Fox. *Trends in Ecology & Evolution*, *28*, 571-572.
- Shelley, A. J., Hernández, L. M., Maia-Herzog, M., Luna Dias, A. P. A., & Garritano, P. R. (2010). *The Blackflies (Diptera: Simuliidae) of Brazil*. In J. R. Arias, S. Golovatch, K. M. Wantzen, & E. Dominguez (Series Eds.). *Aquatic Biodiversity in Latin America* (Vol 6, p. 821). ABLA Series.
- Shepp, D. L., & Cummins, J. D. (1997). Restoration in an urban watershed: Anacostia River of Maryland and the district of Columbia. In J. E. Williams, C. A. Wood, & M. P. Dombeck (Eds.), *Watershed restoration: principles and practices* (pp. 297-317). Bethesda: American Fisheries Society.
- Strieder, M. N. & Corseuil, E. (1992). Atividades de hematofagia em Simuliidae (Diptera, Nematocera) na Picada Verão, Sapiranga, RS-Brasil. *Acta Biologica Leopoldensia*, *14*, 75-98.
- Strieder, M. N., Ronchi, L. H., Stenert, C., Scherer, R. T., & Neiss, U. G. (2006). Medidas biológicas e índices de qualidade da água de uma microbacia com poluição urbana e de curtiúmes no sul do Brasil. *Acta Biologica Leopoldensia*, *28*, 17-24.
- Svensson, J. R., Lindgarth, M., Jonsson, P. R., & Pavia, H. (2012). Disturbance-diversity models: what do they really predict and how are they tested? *Proceedings of the Royal Society: Biological Sciences*, *279*, 2163-2170.
- Ter Braak, C. J. F., & Smilauer, P. (2002). *CANOCO Reference manual and CanoDraw for Windows user's guide: Software for Canonical Community Ordination (version 4.5)*. Wageningen: Bometris.
- Timm, T. (1995). *Ufer- und Auestrukturen und Simuliiden-Plagen*. In C. Steinberg, W. Bernhardt, & H. Klapper (Eds.), *Handbuch Angewandte Limnologie* (p. 28). Weinheim: Kluwer Academic Publishers.
- Travis, B. V., & Vargas, V. M. (1978). Bionomics of black flies (Diptera: Simuliidae) in Costa Rica. Correlations with ecological factors. *Revista Biologia Tropical*, *26*, 335-345.
- Zhang, Y., Malmqvist, B., & Englund, G. (1998). Ecological processes affecting community structure of blackfly larvae in regulated and unregulated rivers: A regional study. *Journal of Applied Ecology*, *35*, 673-686.