

- Schoener, T. W. (1968). The *Anolis* lizards of Bimini: resource partitioning in a complex fauna. *Ecology*, 49, 704-726.
- Sifres, A., Blanca, J., & Nuez, F. (2011). Pattern of genetic variability of *Solanum habrochaites* in its natural area of distribution. *Genetic Resources and Crop Evolution*, 58, 347-360.
- Soberón, J., & Peterson, A. T. (2005). Interpretation of models of fundamental ecological niches and species' distributional areas. *Biodiversity Informatics*, 2, 1-10.
- Vargas, C. D., Rodríguez, E. G., Sánchez, J. J. G., Montes, S. H., Ruiz, A. C., Lápiz, R. I., Puente, P. O., & Martínez, J. L. R. (2005). Adaptación climática de *Lycopersicon* en el occidente de México. En S. Carvajal & E. Pimienta Barrios, *Avances en la Investigación Científica en el CUCBA* (pp. 207-210). Guadalajara, México: Universidad de Guadalajara.
- Yang, Z., Li, Y., Li, P., Zhang, F., & Thomas, B. W. (2016). Effect of difference between day and night temperature on tomato (*Lycopersicon esculentum* Mill.) root activity and low molecular weight organic acid secretion. *Soil Science and Plant Nutrition*, 62, 423-431.
- Yang, X. Q., Kushwaha, S. P. S., Saran, S., Xu, J., & Roy, P. S. (2013). Maxent modeling for predicting the potential distribution of medicinal plant, *Justicia adhatoda* L. in Lesser Himalayan foothills. *Ecological Engineering*, 51, 83-87.
- Zuriaga, E., Blanca, J. M., Cordero, L., Sifres, A., Blas-Cerdán, W. G., Morales, R., & Nuez, F. (2009). Genetic and bioclimatic variation in *Solanum pimpinellifolium*. *Genetic Resources and Crop Evolution*, 56, 39-51.

ANEXO 1

Idoneidad de MNEorigen: los colores oscuros (negro) indican áreas de mayor idoneidad y los claros indican menor idoneidad

APPENDIX 1

MNEorigen suitability: dark colors (black) indicate areas of higher suitability and light colors indicate lower suitability

