



Elizalde, V., García, J.R., Trejo, C., Peña-Valdivia, C.B., Ma. Ybarra, C., & Leyva, O.R. (2021). Seed mass maturity in the terrestrial bromeliad *Hechtia perotensis* (Bromeliaceae), endemic to Mexico. *Revista de Biología Tropical*, 69(3), 843-851. <https://doi.org/10.15517/rbt.v69i3.43477>

<https://doi.org/10.15517/rbt.v69i3.43477>

Seed mass maturity in the terrestrial bromeliad *Hechtia perotensis* (Bromeliaceae), endemic to Mexico


Violeta Elizalde¹;  <https://orcid.org/0000-0002-4801-8782>

José Rodolfo García^{1*};  <https://orcid.org/0000-0002-2086-1073>

Carlos Trejo¹;  <https://orcid.org/0000-0003-3431-4828>

Cecilia Beatriz Peña-Valdivia¹;  <https://orcid.org/0000-0003-4245-0547>

Ma. Carmen Ybarra²;  <https://orcid.org/0000-0002-9634-008X>

Otto Raúl Leyva³;  <https://orcid.org/0000-0002-6150-9367>

1. Postgrado en Botánica, Colegio de Postgraduados, Campus Montecillo, Carretera México-Texcoco, km 35.5, Estado de México, 56230, México; violetaelizalde@gmail.com, garcianr@colpos.mx (*Correspondence), catre@colpos.mx, cecilia@colpos.mx
2. Departamento de Ingeniería Agroindustrial, Universidad Autónoma Chapingo, Carretera México-Texcoco km 38.5, Estado de México, 56230, México; ycydrive@gmail.com
3. Facultad de Ciencias Biológicas y Agropecuarias, Universidad Veracruzana, Camino a Peñuela s/n. C.P.94945 Peñuela, Amatlán de los Reyes, Ver., México; oleyva@uv.mx

Received 13-VIII-2020. Corrected 29-IV-2021. Accepted 03-08-2021.

ABSTRACT

Introduction: *H. perotensis* is a plant with a high potential for ecological restoration because it yields thousands of seeds and grows under low levels of rain, poor soils and contrasting temperatures. However, little is known of the seed mass maturity (high seed germination, low seed fresh weight and low seed moisture content) in this species.

Objective: Assess seed germination in the laboratory of *H. perotensis* during seed development and along the floral stalk (infructescence) in two sites one in rocky location and another near a lake. The hypothesis was that there is a time after flowering in which seeds have highest germination and fresh weight and that the apical, centre and base of the infructescence are different in seed germination and fresh weight in both sites.

Methods: Capsules were collected in two sites one in rocky land (Frijol Colorado, Perote, Veracruz) and another near one lake (Alchichica, Puebla), in the months of August, September and November 2016 and January 2017. A repeated measure design (RMD) was used to analyze the effects of infructescence section on seed weight, moisture content and seed germination (41, 87, 152 and 215 days after flowering). Each evaluation time comprised five replicates, each one with 15 seeds.

Results: Difference in seed germination, seed weight and moisture content between sections of the infructescence was not significant. However, significant differences were found not only between first and last sample dates, but mainly between first and second dates. Eighty-seven days after flower pollination seed moisture content was lower than 20 % and up to 80 % of seed germinated in both sites of sampling.

Conclusions: In this study it was found that the moisture content of *H. perotensis* seed can be used as an indicator of the physiological maturity of the seed and it is also related to germination of the seed.

Key words: ecological restoration; infructescence; moisture content; germination; seed weight.



The genus *Hechtia* Klotzsch is one of the more interesting members of the Mexican Bromeliaceae because it is the only member of the subfamily Hechtioideae and has experienced recent discovery of new species (Ramírez et al., 2014). The estimated number of species in Mexico is 65 (Ramírez & Jiménez, 2012) and 94 % of them are endemic (Ramírez et al., 2014). *Hechtia* plants are terrestrial or lithophile and can grow on volcanic rocks and limestone hills. Bromeliaceae seeds are brown, fusiform, rough, oblong, ellipsoids, one mm wide and three mm long (Espejo et al., 2007). Reyes (2015) suggests that *Hechtia* can be used in ecological restoration programs because it produces a remarkably high quantity of seeds and the ability of plants to grow under low rainfall, on poor soils, and under contrasting temperatures. However, the knowledge of *Hechtia* seed biology is limited (Espejo et al., 2007). Information of seed moisture content, dry weight, and seed germination after seed mass maturity of *H. perotensis* has not been reported. Information about the genus *Hechtia* includes seed morphological descriptions of *H. nususaviorum* (Espejo et al., 2007) and new species descriptions of *H. schottii* (Escobedo, 2012), *H. santanae* (Ramírez et al., 2016), *H. flexilifolia*, *H. huamelulaensis*, *H. nivea* (Ramírez et al., 2014).

Seed germination characteristics are fundamental for species conservation (Baskin & Baskin, 1998). In consequence, it is necessary to know the timing of seed harvest as it influences seed quality (physical and physiological) and storage effects (Willan, 1985) to initiate nursery programs and store seeds in germplasm banks. Seed germination is sensitive to environmental conditions (Leck et al., 2008). Sankhla and Chawan (1980) found that seed moisture content is important in regulating seed germination. For example, *Phaseolus trilobus* seeds had 100 % seed germination at 30 % moisture content and 0 % with moisture content of 2 %. In addition, seed quality continues to improve during development and even after maximum dry weight (Pieta & Ellis, 1991) as seen with seeds of beans (Sanhewe & Ellis,

1996) and cereals (Hay & Probert, 1995). Seed development of *H. perotensis* has not been reported and this, together with timing of seed harvest is a first requirement to start propagation in nurseries and to store seeds with maximum seed quality in seed banks. The objective of this research was to assess seed moisture content, seed weight and seed germination of *H. perotensis* during seed development in two sites one in a rocky location and another near a lake. The hypothesis was that after flowering there is a time nearly at the end of seed mass maturity where seeds have low weight due to seed drying, low moisture content and maximum germination. Also, that seed quality differs among three longitudinal sections of the infructescence due to differences in maturity along the stalk.

MATERIAL AND METHODS

Sample sites: *H. perotensis* seeds were collected at site 1 a rocky land, Frijol Colorado, Perote, Veracruz (2 416 m.a.s.l, 19°32'13.95" N & 97°22'98" W) and site 2, Tepeyahualco at the lake of Alchichica, Puebla, México (2 326 m.a.s.l, 19°25'11" N & 97°23'56" W) (geolocation with Garmin eTrex 10 & Google Earth®) (Google Earth, n.d.-a; Google Earth, n.d.-b). The sites were selected to investigate the effect of the lake (site 2) in the relative humidity and on seed mass maturity compared with site 1 plants growing in rocky land. The climate of each region is semiarid with warm summers BS1kw(i'gw)" (García, 1988). The vegetation is typical of a xeric shrublands (Ramírez et al., 2014). Soil was alkaline, not saline, loamy and poor in nutrients in both sites (laboratory analysis).

Plant selection: In a first visit in June 2016 thereafter called day 0, male and female plants in both sites had flowers and insect pollinations. After one month, plants of both sites had green immature capsules and at this moment a white perforated sleeve shape cloth with diameter of 1 mm was inserted along the infructescence in order to keep the seeds

inside the cloth in case of dehiscence. Each infructescence length was registered with a measuring tape, divided in three and adjusting a lace in each section to keep the mature seeds separated.

Capsule sampling: Capsules of *H. perotensis* in both sites were collected in August, September, November 2016 and January 2017. The mean temperature was 15.3 and 17.6 °C, and mean relative humidity for 56.8 and 60 % (data logger Hobo), of site 1 and 2, respectively during the sample months. Temperature and relative humidity were measured to compare the environment characteristics.

Once capsule colour changed from green to brown around 41 d after flowering (DAF) 25 capsules were harvested from each of the base, centre, and apex of the infructescence. Two to four capsule sample dates were made according to the availability of capsules. Samples were made at 41, 87, 152 and 215 DAF in plants named A in site 1 and B in site 2. Plant named C and D of site 1 and 2 respectively were

sampled at 41 and 87 DAF due to less availability of capsules.

Seeds extracted from capsules with fine clippers were stored at (25 ± 1 °C) in seal aluminium bags. Seed weight, humidity and germination evaluations were made within one week after each harvest period.

Seed weight: Seed weight was determined in 225 seeds by infructescence in a precision balance (± 0.0001 g; Scientech SA 120) taking 75 seeds from the apex, centre and base of the infructescence this procedure was repeated for each sample date.

Seed moisture content: Seed fresh weight of five replicates each with 15 seeds was determined from the apex, centre, and base of the infructescence this procedure was repeated for each sample date. Seeds were kept at 50 ± 1 °C in an oven until they reach constant weight (5 d). Seed moisture content was calculated using the following expression (International Seed Testing Association, 2010).

$$\text{Seed moisture content} = \frac{\text{Seed fresh weight} - \text{Seed dry weight}}{\text{Seed fresh weight}} \times 100$$

Germination: Seeds were placed in petri dishes with filter paper on the bottom and 10 mL of distilled water was added. Five replicates with fifteen seeds (per petri dish of 14 cm) were tested from the apex, centre, and base of the infructescence this procedure was repeated for each sample date. Seeds were superficially disinfested by immersion in sodium hypochlorite (60 g/L of Cl active for 1 min) 1 % (v:v in water). Petri dishes were placed in an incubator (Thermo Scientific. USA: model 846) set at 25 ± 1 °C and photoperiod of 12 h darkness x 12 h light (12 μmol m⁻² s⁻¹ of irradiation). A seed was considered germinated when 1 mm of radicle was exposed using a digital caliper Vernier. Evaluations were made at 8 and 15 d from placement in the incubator (International Seed Testing Association, 2010).

Seeds in development: Seeds were observed with a stereo microscope (Leica EZ4 HD) to identify embryo development this procedure was repeated for each sample date. Seed testa was dissected near the embryo and the embryo was extracted with tweezers (International Seed Testing Association, 2010).

Experimental design: A repeated measures design (RMD) was used to analyze the effects of time (day 41, 87, 152 and 215) and infructescence section on seed weight, moisture content, seed germination and seed development at each site. Each evaluation time comprised five replicates each of 15 seeds as an experimental unit from the apex, centre, and base of the infructescence. Multiple




Fig. 1. A. B. C. and D. Germination, seed moisture content, seed fresh weight and seeds in development from the plant A site 1 (Frijol Colorado, Perote, Veracruz) and **E. F. G. and H** of plant B, from site 2 (Tepeyahualco at the lake of Alchichica, Puebla). Each column represents the average of 5 replicates of 15 seeds each. The apex (■), centre (▒) and base (▨) of the stalk is represented by three columns. The letters on the columns represent the effects of the time in each section of the stalk by date of days after flowering.

comparisons were made with the general lineal model ($\alpha = 0.05$).

RESULTS

Plant A site 1 Frijol Colorado, Perote, Veracruz: Seeds of the three sections of the stalk (base, medium and apex) were similar on

germination, moisture content and seed fresh weight (Fig. 1A, Fig. 1B, Fig. 1C) in each date sampled ($P > 0.05$). However, there was statistical difference over time; at 41 DAF seed fresh weight average at three sections was 0.036 g, moisture content of 76 % and seed germination of 20 %. Whereas seeds sampled at 215 DAF had seed fresh weight average at the three

sections of 0.0139 g, seed moisture content of 6 % and 94 % of seed germination ($P \leq 0.05$). Seeds in development were similar in the three sections of the stalk at 41 DAF with an average of 80 % of seeds. At 87, 152 and 215 DAF seeds in development were on average 6 % without differences in the sections of the stalk (Fig. 1D).

Plant B site 2 Tepeyahualco at the lake of Alchichica, Puebla: Seed germination, moisture content and seed fresh weight (Fig. 1E, Fig. 1F, Fig. 1G) was similar in the three sections of the stalk in each date sampled, however seed germination, moisture content and seed fresh weight showed evident differences with time. On average, seed germination at 215 DAF (98 %) was 2.6 times higher than that at 41 DAF (37 %) (Fig. 1E). Conversely, seed moisture content diminished from 69 to 6 %, nearly ten times in the same period (Fig. 1F).

Seed fresh weight (Fig. 1G) was different ($P \leq 0.05$) in the three sections of the stalk over time. The average seed fresh weight was more than double at 41 DAF (0.036 g) compared with 215 DAF (0.016 g) this decrease of seed moisture was due to seed drying.

Seeds in development were similar in the three sections of the stalk at 41 DAF with an average of 60 % of seeds. At 87, 152 and 215 DAF seeds in development were on average 4 % without differences in the sections of the stalk (Fig. 1H).

Plant C site 1 Frijol Colorado, Perote, Veracruz: Seeds of the three sections of the stalk did not differ in germination, moisture content and seed fresh weight at 41 and 87 DAF (Fig. 2A, Fig. 2B, Fig. 2C) ($P > 0.05$). However, the average seed germination at 41 and 87 DAF was 23 and 95 % respectively (Fig. 2A). Seed moisture content and seed fresh weight of the three sections on average was 72 % and 0.035 g at 41 DAF. At 87 DAF seed moisture content diminished to 29 % and seed fresh weight was 0.017 g (Fig. 2B, Fig. 2C).

Seeds in development were similar in the three sections of the stalk at 41 DAF with an average of 76 % of seeds. At 87 DAF seeds

in development were on average 4 % without differences between the sections of the stalk (Fig. 2D).

Plant D site 2 Tepeyahualco at the lake of Alchichica, Puebla: Seed germination, moisture content and seed fresh weight (Fig. 2E, Fig. 2F, Fig. 2G) was similar in both dates sampled. Also, seed moisture content and seed fresh weight differed over time ($P \leq 0.05$), but not between stalk sections. The average seed germination of the stalk was 83 % at 87 DAF (Fig. 2E). Seed moisture content and fresh weight (Fig. 2F, Fig. 2G) decreased to 9 % and 0.0144 g respectively averaged across, the three sections of the stalk at 87 DAF.

Seeds in development were similar in the three sections of the stalk at 41 DAF with an average of 35 % of seeds. At 87 DAF seeds in development were on average 15 % differing between the apex and centre or base of the stalk (Fig. 2H).

DISCUSSION

Our research showed homogeneity and high seed germination (87 %) of *H. perotensis* at 87 DAF at the two sites of seed collection and across three sections of the infructescence. In previous research Elizalde et al., (2017) showed variation of seed germination of *H. perotensis* from two different harvest dates (20 % in 2012 and 92 % in 2015), and as a consequence, it was necessary to establish the time at which *H. perotensis* reaches high seed quality.

Gutterman (1980) confirmed that seed position in the fruit or infructescence is important for seed germination. The perennial shrub *Mesembryanthemum nodiflorum* had different seed germination according to the position of the infructescence in which the seed grows. For example, in the apex seed germination was of 61 %, in the centre 5.5 % and in the base 1 %. *M. nodiflorum* grows in the desert and the difference in germination of the fruit is related with the cycle life of this plant. As a consequence, undetermined factors could have influenced seed quality along an infructescence,




Fig. 2. A. B. C. and D. Germination, seed moisture content, seed fresh weight and seeds in development from the plant C, site 1 (Frijol Colorado, Perote, Veracruz) and **E. F. G. and H** of plant D from site 2 (Tepeyahualco at the lake of Alchichica, Puebla). Each column represents the average of 5 replicates of 15 seeds each. The apex (■), centre (▒) and base (▨) of the stalk is represented by three columns. The letters on the columns represent the effects of the time in each section of the stalk by date of days after flowering.

such as environmental conditions during the development of the mother plant, flower aperture, and occurrence of pollination insects (Gray & Thomas, 1982).

In contrast, Gesch et al., (2016) did not find differences in seed germination between different seed positions in plants of *Thlaspi arvense* L. Similarly, in our research seeds of

H. perotensis did not show such locational differences in germination.

Willson and Price (1977) found that production of more flowers in one plant attracts more pollinating insects and as a result plant produce more seeds. McKinney et al., (2012) showed that flowering timing coincides with activity of pollinating insects to improve

homogenous flower pollination (Hegland et al., 2009). *H. perotensis* was flowering with pollination insects and we assumed that flower pollination was efficient. In addition, the capsule formation was homogenous in the three sections of the stalk mainly in seed germination.

Seed moisture content diminished gradually during seed development: the reserves are accumulated (sugars) and solutes (proline) in the vacuole and there is water displacement to protect cell membranes. After, physiological seed maturity (mass maturity) orthodox seeds increase the process of seed drying to reach the end of seed maturation (Bewley et al., 2013). We found that seeds of *H. perotensis* improved seed germination by about 80 % when seed moisture content decrease to 20 % at the last sample analyzed. *H. perotensis* seeds drying in an environment with low relative humidity (60 %) and a medium temperature 16 °C, reaching a low moisture content in about 45 days.

Bhawna et al., (2011) indicated that seed maturity of *Prunus cerasoides* was at seed moisture content of 31.96 % ± 1.42 %. According to Mai-Hong et al., (2003) 100 % seed germination of *Peltophorum pterocarpum* (DC) K. Heyne is reach when seed moisture content was 56 % with maximum seed dry weight of 0.5 g. The seed moisture content has been used as a reliable indicator of seed maturity in several studies e.g., by Shah et al., (2010). In our study we found that seed moisture is an indicator of seed maturity in *H. perotensis*. Harvest seeds at 41 DAF were in development with an average moisture content of 70 %. Seeds reached embryo maturity and increased germination at 87 DAF with only 10 % of seeds in development. At this stage, maturation is approaching and maturation drying is the event for orthodox seeds to enter in a metabolic quiescent state. Seeds can remain in this state for days to years with high viability (Bewley et al., 2013).

Shah et al., (2006) showed that seed moisture content of 23.4 to 36.1 % can be associated with an optimum seed germination of *Pyracantha crenulata*. Bewley et al., (2013) reported that seed physiological maturity (vigour and seed germination) is reached when seed dry

weight is greatest, and after this moment seeds start to decrease in quality. For example, in soya seeds maximum dry weight was obtained 45 d after flowering. For seeds of *Caesalpinia lutea* and *C. wrightii* physiological maturity was reached around 18 to 21 d after flowering (Kaliangile & Grabe, 1988). Our results agree with the above authors and demonstrated that *H. perotensis* reach minimum seed fresh weight (low seed moisture content) with coincidence of maximum seed germination at the end of the samples periods (87 DAF).

Maturity of seeds also has been associated with seed moisture content, for example, *Ricinus communis* L. seed maturity was reported to occur at 22 % seed moisture content (Vallejos et al., 2011). Seeds of *Aesculus indica* Colebr, *Albizia lebbek* and *Celtis australis* had physiological maturity at 58, 52 and 32 % seed moisture content respectively (Majeed et al., 2010; Bhardwaj et al., 2002). Our research showed that seeds of *H. Perotensis* reached physiological maturity with fresh seed weight of 0.010 to 0.015 g, moisture content lower than 20 % and seed germination of 85 %. Similar results of seed fresh weight were reported by Elizalde et al., (2017), however they mainly focused on methods to improve seed germination.

Sankhla and Chawan (1980) claim that seeds with high moisture content could produce low seed germination. The authors explained that a seed before drying is in development with immature embryos and in the process of storage of nutritive reserves. Therefore, seed drying is necessary to reach high germination percentages and seed maturity. After seed drying the events of seed development stop and with seed water uptake the seed germination metabolism starts (Bewley et al., 2013).

We conclude that *H. perotensis* reached seed mass maturity 87 d after flowering with 20 % moisture content. With this knowledge the point of harvest of seeds can be used to store seeds with high percentage of germination in seed banks and to start a plant nursery and use *H. perotensis* for ecological restoration in their origin place and has the potential to use this plant in other regions with similar



environments. Both sites studied reached high seed germination at the same time.

Ethical statement: authors declare that they all agree with this publication and made significant contributions; that there is no conflict of interest of any kind; and that we followed all pertinent ethical and legal procedures and requirements. All financial sources are fully and clearly stated in the acknowledgements section. A signed document has been filed in the journal archives.

ACKNOWLEDGMENTS

To the Consejo Nacional de Ciencia y Tecnología (CONACYT) for the scholarship for the first author. Many thanks to the reviewers and editor that help to improve the manuscript.

RESUMEN

Madurez de las semillas en la bromelia terrestre *Hechtia perotensis* (Bromeliaceae), endémica de México

Introducción: *H. perotensis* es una planta con un alto potencial para la restauración ecológica debido a que produce miles de semillas y crece con niveles bajos de lluvia, suelos pobres y temperaturas contrastantes. Sin embargo, el desarrollo de la madurez de las semillas (germinación alta, baja humedad y bajo peso fresco de semilla) en esta especie es muy poco conocido.

Objetivo: Evaluar la germinación en el laboratorio de las semillas de *H. perotensis*, durante el desarrollo de la semilla y a lo largo del tallo floral (infrutescencia) en dos sitios; uno en un área rocosa y otro cerca de un lago. La hipótesis fue que hay un tiempo después de la floración, en el que las semillas presentan la mayor germinación y menor peso fresco, y que las secciones apical, central y base de la infrutescencia, la germinación y el peso fresco de la semilla son diferentes en los dos sitios.

Métodos: se recolectaron las cápsulas en dos sitios uno en un área rocosa (Frijol Colorado, Perote, Veracruz) y otro cerca de un lago (Alchichica, Puebla), en los meses de Agosto, Septiembre y Noviembre de 2016 y Enero de 2017. Se usó un diseño de medidas repetidas (DMR) para analizar los efectos de la sección de la infrutescencia sobre el peso fresco de la semilla, el contenido de humedad y la germinación de la semilla (41, 87, 152 y 215 días, después de la floración). Cada tiempo de evaluación comprendió cinco réplicas y 15 semillas.

Resultados: La diferencia en la germinación de la semilla, el peso de la semilla y el contenido de humedad entre las

secciones de la infrutescencia no fue significativa. Sin embargo, se encontraron diferencias significativas entre las fechas de muestreo inicial y final, pero sobre todo entre la primera y la segunda fecha. Asimismo, ochenta y siete días después de la polinización de las flores, el contenido de humedad de las semillas fue inferior al 20 % y superior al 80 % en la germinación de las semillas en ambos sitios de muestreo.

Conclusiones: en este estudio se encontró que el contenido de humedad de la semilla de *H. perotensis*, puede usarse como un indicador de la madurez fisiológica de la semilla y también está relacionado con la germinación de la semilla.

Palabras clave: restauración ecológica; infrutescencia; contenido de humedad; germinación; peso de semilla.

REFERENCES

- Baskin, C., & Baskin, J. (1998). *Seeds: ecology, biogeography, and evolution of dormancy and germination*. Academic Press.
- Bewley, J., Bradford, K., Hilhorst, H., & Nonogaki, H. (2013). *Physiology of development, germination and dormancy*. Springer Science Business Media.
- Bhardwaj, S., Panwar, P., & Kumar, M. (2002). Physical and biochemical changes in seeds as maturity index for harvesting *Albizia lebbek* seeds. *Journal of Hill Research*, 15(1), 52–55.
- Bhawna, T., Ashish, T., Shruti, S., & Neerja, P. (2011). Physical attributes as indicator of seed maturity and germination enhancement in Himalayan Wild Cherry (*Prunus cerasoides* D. Don.). *New Forests*, 41, 139–146.
- Elizalde, V., García, J., Peña-Valdivia, C., Ybarra, C., Leyva, O., & Trejo, C. (2017). Viabilidad y germinación de semillas de *Hechtia perotensis* (Bromeliaceae). [Seed viability and germination of *Hechtia* (Bromeliaceae)]. *Revista de Biología Tropical*, 65(1), 153-165.
- Escobedo, S. (2012). *Mecanismos de dispersión de semillas en las Bromelias. Desde el herbario CICY 4: 22-23*. http://www.cicy.mx/sitios/desde_herbario/
- Espejo, S., López, F., Ramirez, M., & Martínez, C. (2007). Dos nuevas especies de *Hechtia* (Bromeliaceae) de México. *Acta Botánica Mexicana*, 78, 97–109.
- García, E. (1988). *Modificaciones al sistema de clasificación climática de Köppen*. Universidad Nacional Autónoma de México.
- Gesch, R., Royo-Esnal, A., Edo-Tena, E., Recasens, J., Isbell, T., & Forcella, F. (2016). Growth environment but not seed position on the parent plant affect seed germination of two *Thlaspi arvense* L. populations. *Industrial Crops and Products*, 84, 241–247.

- Google Earth. (n.d.-a). *Frijol Colorado, Perote, Veracruz*. Recover on July 14, 2021 from https://earth.google.com/web/search/frijol+colorado/@19.55318431,-97.37798852,2433.58431515a,10930.6604354d,34.99998921y,0h,0t,0r/data=CigiJgokCYtftwEzbTNAEZ9j_ux9aT-NAGTSh-p_gWFjAISX4BY4tWljA
- Google Earth. (n.d.-b). *Alchichica, Puebla*. Recover on July 14, 2021 from <https://earth.google.com/web/search/alchichica+puebla/@19.41931892,-97.39887023,2323.53443059a,681.20123398d,30.00019y,0h,0t,0r/data=CigiJgokCZeeWHSnjNAEb7OAbWWjzNAGbzLrRTaUljAIExtrkCwWljA>
- Gray, D., & Thomas, T. (1982). Seed germination and seedling emergence as influenced by the position of development of the seed on, and chemical applications to, the parent plant. In A. A. Khan (Ed.), *The Physiology and Biochemistry of Seed Development, Dormancy and Germination* (pp. 81–110). Elsevier Biochemical Press.
- Gutterman, Y. (1980). Annual rhythm and position effect in the germinability of *Mesymbrianthemum nodiflorum*. *Israel Journal of Botany*, 29, 93–97.
- Hay, F., & Probert, R. (1995). The effect of different drying conditions and maturity on desiccation tolerance and seed longevity in *Digitalis purpurea* L. *Annals of Botany*, 76, 639–647.
- Hegland, S., Nielsen, A., Lazaro, A., Bjerknes, A., & Totland, O. (2009). How does climate warming affect plant-pollinator interactions? *Ecology Letters*, 12, 184–195.
- International Seed Testing Association (2010). *International Rules for Seed Testing*. International Seed Testing Association.
- Kaliangile, I., & Grabe, D. (1988). Seed maturation in *Cuphea*. *Journal of Seed Technology*, 12(2), 107–113.
- Leck, M., Parker, V., & Simpson, R. (2008). *Seedling ecology and evolution*. Cambridge University Press.
- Mai-Hong, T., Hong, T.D., Hien, N.T., & Ellis, R.H. (2003). Onset of germinability, desiccation tolerance and hardseededness in developing seeds of *Peltophorum pterocarpum* (DC) k. Heyne (Caesalpinioideae). *Seed Science Research*, 13, 323–327.
- Majeed, M., Khan, M., Bashir, A., & Qaisar, K. (2010). Maturity indices of Indian horse chestnut (*Aesculus indica* Colebr.) seeds under temperate Kashmir conditions. *Forestry Studies in China*, 12(1), 45–48.
- McKinney, A., CaraDonna, P., Inouye, D., Barr, B., Bertelsen, C., & Waser, N. (2012). Asynchronous changes in phenology of migrating Broad-tailed Hummingbirds and their early-season nectar resources. *Ecology*, 93, 1987–1993.
- Pieta, F., & Ellis, R. (1991). The development of seed quality in spring barley in four environments. I. Germination and longevity. *Seed Science Research*, 1, 163–177.
- Ramírez, M., & Jiménez, C. (2012). A new species of *Hechtia* (Hechtioideae: Bromeliaceae) from Puebla, Mexico. *Phytotaxa*, 42, 1–8.
- Ramírez, M., Jiménez, F., Fernández-Concha, G., & Pinzón, J. (2014). Three new species and growth patterns in *Hechtia* (Bromeliaceae: Hechtioideae). *Phytotaxa*, 178, 113–127.
- Ramírez, M., Carrillo, R., Tapia, M., & Cetzal, I. (2016). An addition to genus *Hechtia* (Hechtioideae; Bromeliaceae) from Jalisco, Mexico. *Phytotaxa*, 266(4), 261–270.
- Reyes, S. (2015). *Conservación y restauración de cactáceas y otras plantas suculentas mexicanas*. Comisión Nacional Forestal.
- Sanhewe, A., & Ellis, R. (1996). Seed development and maturation in *Phaseolus vulgaris*. II. Post-harvest longevity in air-dry storage. *Journal of Experimental Botany*, 47, 959–965.
- Sankhla, R., & Chawan, D. (1980). Effect of different seed moisture levels on the germination behaviour of *Phaseolus trilobus*. *Biologia Plantarum*, 22(5), 388–391.
- Shah, S., Tewari, A., & Bisht, S. (2006). Seed maturation indicators in *Pyraecantha crenulata* Roxb. in Kumaun central Himalaya. *New Forests*, 32, 1–7.
- Shah, S., Tewari, A., Tewari, B., & Singh, R. (2010). Seed maturity indicators in *Myrica esculenta*, Buch-Ham. Ex. D. Don.: a multipurpose tree species of subtropical temperate Himalayan region. *New Forests*, 40, 9–18.
- Vallejos, M., Rondanini, D., & Wassner, F. (2011). Water relationships of castor bean (*Ricinus communis* L.) seeds related to final seed dry weight and physiological maturity. *European Journal of Agronomy*, 35, 93–101.
- Willan, R. (1985). *A guide to forest seed handling*. FAO Forestry paper 20/2. FAO.
- Willson, M., & Price, P. (1977). The evolution of inflorescence size in *Asclepius* (Asclepiadaceae). *Evolution*, 31, 495–511.